

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 674 111 A1

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:

28.06.2006 Bulletin 2006/26

(21) Application number: **05760156.9**

(22) Date of filing: **08.07.2005**

(51) Int Cl.:

A61K 39/395 ^(2006.01) **A61P 1/16** ^(2006.01)
A61P 35/00 ^(2006.01) **C07K 14/47** ^(2006.01)
C07K 16/18 ^(2006.01) **C12N 15/09** ^(2006.01)
C12P 21/08 ^(2006.01)

(86) International application number:
PCT/JP2005/013103

(87) International publication number:
WO 2006/006693 (19.01.2006 Gazette 2003/06)

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI
SK TR**

Designated Extension States:

AL BA HR MK YU

(30) Priority: **09.07.2004 JP 203637**

(71) Applicant: **CHUGAI SEIYAKU KABUSHIKI KAISHA
Tokyo, 115-8543 (JP)**

(72) Inventors:

- **NAKANO, Kiyotaka,**
c/o Chugai Seiyaki K.K.
Nihari-gun,
Ibaraki 3004101 (JP)
- **YOSHINO, Takeshi,**
c/o Chugai Seiyaki K.K.
Nihari-gun,
Ibaraki 3004101 (JP)
- **NEZU, Jun-ichi,**
c/o Chugai Seiyaki K.K.
Nihari-gun,
Ibaraki 3004101 (JP)
- **TSUNODA, Hiroyuki,**
c/o Chugai Seiyaki K.K.
Nihari-gun,
Ibaraki 3004101 (JP)

- **IGAWA, Tomoyuki**
Gotemba-shi,
Shizuoka 4128513 (JP)
- **KONISHI, Hiroko**
Gotemba-shi,
Shizuoka 4128513 (JP)
- **TANAKA, Megumi**
Gotemba-shi,
Shizuoka 4128513 (JP)
- **SUGO, Izumi**
Gotemba-shi,
Shizuoka 4128513 (JP)
- **KAWAI, Shigeto**
Kamakura-shi,
Kanagawa 2478530 (JP)
- **ISHIGURO, Takahiro**
Kamakura-shi,
Kanagawa 2478530 (JP)
- **KINOSHITA, Yasuko**
Kamakura-shi,
Kanagawa 2478530 (JP)

(74) Representative: **Woods, Geoffrey Corlett**
J.A. KEMP & CO.
Gray's Inn
14 South Square
London WC1R 5JJ (GB)

(54) **ANTI-GLYPICAN 3 ANTIBODY**

(57) An antibody capable of binding to a specific region of glypican 3, as well as a humanized antibody created based on that antibody are disclosed. The anti-GPC3 antibody of the invention has a higher ADCC activity and CDC activity compared with those of a conven-

tional antibody. The antibody of the present invention is useful as a cell growth inhibitor, an anticancer agent and an agent for diagnosis of cancers.

EP 1 674 111 A1




FIG.10

Description**BACKGROUND OF THE INVENTION**5 **Field of the Invention**

[0001] The present invention relates to an anti-glypican 3 antibody, a cell growth inhibitor and an anticancer agent containing the antibody as an active ingredient.

10 **Description of Related Art**

[0002] Glypican 3 (GPC3) is one of the glypican family of heparan sulfate proteoglycans that are present on cell surfaces. It is suggested that GPC3 may be involved in cell division in development or cancer cell growth, however, its function has not been well elucidated yet.

15 **[0003]** It has been found that a certain type of antibody binding to GPC3 has a cell growth-inhibiting activity via an antibody-dependent cell-mediated cytotoxicity (ADCC) activity and a complement-dependent cytotoxicity (CDC) activity (International Patent Application WO 2003/000883). In addition, it has been suggested that GPC3 is cleaved in vivo and secreted into blood as a secreted form of GPC3, and the diagnosis of cancers may be possible by using an antibody capable of detecting the secreted form of GPC3 (International Patent Applications WO 2004/022739, WO 03/100429 and WO 2004/018667).

20 **[0004]** When developing an anticancer agent based on the cytotoxicity activity of an antibody, it is preferred that the antibody to be used has high ADCC activity or CDC activity. Accordingly, an anti-GPC3 antibody having a high cytotoxicity activity has been desired as an antibody recognizing GPC3.

25 **[0005]** An object of the present invention is to provide an anti-GPC3 antibody having a higher ADCC activity and CDC activity compared with those of a conventional antibody.

SUMMARY OF THE INVENTION

30 **[0006]** The present inventors have succeeded in obtaining an antibody having a higher cytotoxicity activity compared with that of a conventional anti-glypican 3 antibody. Furthermore, they analyzed epitopes for such an antibody and succeeded in determining the regions on GPC 3 recognized by the antibody with a high cytotoxicity activity.

[0007] In one aspect, the present invention provides an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 of any one of (1) - (12):

- 35 (1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 123, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 124, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 125;
- (2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 109, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 110, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 111;
- 40 (3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 106, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 107, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 108;
- (4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 132, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 133, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 134;
- (5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 106, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 135, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 136;
- 45 (6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 126, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 127, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 128;
- (7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 129, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 130, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 131;
- (8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 103, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 104, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 105;
- 50 (9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 118, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 121, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 122;
- (10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 115, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 116, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 117;
- 55 (11) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 112, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 113, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 114; or
- (12) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 118, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 119, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 120.

[0008] In another aspect, the invention provides an antibody comprising a light chain variable region having CDRs 1, 2 and 3 of any one of (1) - (13):

- (1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 143, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
- (2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 143, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 145;
- (3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 140, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 141, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 142;
- (4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 167, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 168, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 169;
- (5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 170, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 171;
- (6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 159, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 160, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 161;
- (7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 162, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 147, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 163;
- (8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 164, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 165, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 166;
- (9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 137, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 138, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 139;
- (10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 155, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 156, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 157;
- (11) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 149, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 150, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 151;
- (12) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 146, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 147, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 148; or
- (13) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 152, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 153, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 154.

[0009] Preferably, the antibody of the invention is selected from the group consisting of any one of (1) - (13):

- (1) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 143, 144 and 158, respectively;
- (2) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 109, 110 and 111, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 143, 144 and 145, respectively;
- (3) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 106, 107 and 108, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 140, 141 and 142, respectively;
- (4) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 132, 133 and 134, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 167, 168 and 169, respectively;
- (5) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 106, 135 and 136, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 170, 144 and 171, respectively;
- (6) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 126, 127 and 128, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 159, 160 and 161, respectively;
- (7) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 129, 130 and 131, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 162, 147 and 163, respectively;
- (8) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 129, 130 and 131, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 164, 165 and 166, respectively;
- (9) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 103, 104 and 105, respectively, and a light chain variable region having CDRs

1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 137, 138 and 139, respectively;
 (10) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 118, 121 and 122, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 155, 156 and 157, respectively;
 (11) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 115, 116 and 117, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 149, 150 and 151, respectively;
 (12) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 112, 113 and 114, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 146, 147 and 148, respectively; and
 (13) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 118, 119 and 120, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 152, 153 and 154, respectively.

[0010] In another aspect, the invention provides an antibody having a heavy chain variable region of any one of (1) - (7):

- (1) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84;
- (2) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85;
- (3) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86;
- (4) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87;
- (5) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 88;
- (6) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 89; or
- (7) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 90.

[0011] In another aspect, the invention provides an antibody having a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92.

[0012] Preferably, the antibody of the invention is selected from the group consisting of the antibody of any one of (1) - (7):

- (1) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (2) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (3) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (4) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (5) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 88 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (6) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 89 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92; and
- (7) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 90 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92.

[0013] In another aspect, the invention provides an antibody comprising a light chain variable region having CDRs 1, 2 and 3 of any one of (1) - (15):

- (1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 174, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
- (2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 175, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
- (3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 176, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
- (4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 177, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
- (5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 178, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
- (6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 179, CDR2 comprising the amino acid

sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 180, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 181, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 182, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 183, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (11) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 184, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (12) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 185, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (13) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 186, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; (14) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 187, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158; or (15) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 188, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158.

[0014] In another aspect, the invention provides an antibody selected from the group consisting of the antibody of (1) - (15):

[illegible]

1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 185, 144, and 158, respectively;
 (13) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124, and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 186, 144 and 158, respectively;
 (14) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 187, 144 and 158, respectively; and
 (15) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 188, 144 and 158, respectively.

[0015] In further aspect, the invention provides an antibody having a light chain variable region selected from (1) - (15):

- (1) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 191;
- (2) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 192;
- (3) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193;
- (4) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 194;
- (5) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 195;
- (6) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 196;
- (7) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 197;
- (8) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 198;
- (9) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 199;
- (10) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 200;
- (11) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 201;
- (12) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 202;
- (13) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 203;
- (14) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 204; and
- (15) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 205.

[0016] In another aspect, the invention provides an antibody having a light chain variable region selected from the group consisting of (1) - (15):

- (1) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 191;
- (2) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 192;
- (3) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193;
- (4) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 194;
- (5) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 195;
- (6) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 196;
- (7) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 197;
- (8) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 198;
- (9) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 199;
- (10) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 200;
- (11) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 201;
- (12) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 202;
- (13) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 203;
- (14) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 204; and
- (15) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 205; and a heavy chain variable region selected from the group consisting of (1) - (7):

- (1) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84;
- (2) a heavy chain variable region comprising; the amino acid sequence set forth in SEQ ID NO: 85;
- (3) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86;
- (4) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87;
- (5) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 88;
- (6) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 89; and
- (7) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 90.

[0017] The heavy chain variable region, a light chain variable region, and the amino acid sequence of the CDRs 1, 2

EP 1 674 111 A1

and 3, as well as the SEQ ID NOs are summarized in the table below.

5

10

15

20

25

30

35

40

45

50

55

Antibody and variable regions		SEQ ID NO
M3C11	H	22
M13B3	H	23
M1E7	H	24
M3B8	H	25
M11F1	H	26
M19B11	H	27
M6B1	H	28
M18D4	H	29
M5B9	H	30
M10D2	H	31
L9G11	H	32
M3C11	L	44
M13B3	L	45
M1E7	L	46
M3B8	L	47
M11F1	L	48
M19B11	L	49
M6B1	L	50
M18D4	L	51
M5B9	L	52
M10D2	L	53
L9G11	L	54
GC199	H	60
GC202	H	61
GC33	H	62
GC179	H	63
GC194	H	64
GC199	L	71
GC202	L	72
GC33	L	73
GC179	L	74
GC194(1)	L	75
GC194(2)	L	76
GC33.ver.a	H	84
GC33.ver.c	H	85
GC33.ver.f	H	86
GC33.ver.h	H	87

EP 1 674 111 A1

Table continued

5

10

15

20

25

30

35

40

45

50

55

Antibody and variable regions		SEQ ID NO
GC33.ver.i	H	88
GC33.ver.j	H	89
GC33.ver.k	H	90
GC33.ver.a	L	92
M13B3(H)	CDR1	103
	CDR2	104
	CDR3	105
M3B8(H)	CDR1	106
	CDR2	107
	CDR3	108
M11F1(H)	CDR1	109
	CDR2	110
	CDR3	111
M5B9(H)	CDR1	112
	CDR2	113
	CDR3	114
M6B1(H)	CDR1	115
	CDR2	116
	CDR3	117
M10D2(H)	CDR1	118
	CDR2	119
	CDR3	120
L9G11(H)	CDR1	118
	CDR2	121
	CDR3	122
GC33(H)	CDR1	123
	CDR2	124
	CDR3	125
GC179(H)	CDR1	126
	CDR2	127
	CDR3	128
GC194(H)	CDR1	129
	CDR2	130
	CDR3	131
GC199(H)	CDR1	132
	CDR2	133
	CDR3	134
GC202(H)	CDR1	106

EP 1 674 111 A1

Table continued

Antibody and variable regions		SEQ ID NO
	CDR2	135
	CDR3	136
M13B3(L)	CDR1	137
	CDR2	138
	CDR3	139
M3B8(L)	CDR1	140
	CDR2	141
	CDR3	142
M11F1(L)	CDR1	143
	CDR2	144
	CDR3	145
M5B9(L)	CDR1	146
	CDR2	147
	CDR3	148
M6B1(L)	CDR1	149
	CDR2	150
	CDR3	151
M10D2(L)	CDR1	152
	CDR2	153
	CDR3	154
L9G11(L)	CDR1	155
	CDR2	156
	CDR3	157
GC33(L)	CDR1	143
	CDR2	144
	CDR3	158
GC179(L)	CDR1	159
	CDR2	160
	CDR3	161
GC194(L)1	CDR1	162
	CDR2	147
	CDR3	163
GC194(L)2	CDR1	164
	CDR2	165
	CDR3	166
GC199(L)	CDR1	167
	CDR2	168
	CDR3	169

Table continued

Antibody and variable regions		SEQ ID NO
GC202(L)	CDR1	170
	CDR2	144
	CDR3	171
GC33(L)	G34A	174
GC33(L)	G34D	175
GC33(L)	G34E	176
GC33(L)	G34F	177
GC33(L)	G34H	178
GC33(L)	G34N	179
GC33(L)	G34P	180
GC33(L)	G34Q	181
GC33(L)	G34I	182
GC33(L)	G34K	183
GC33(L)	G34L	184
GC33(L)	G34V	185
GC33(L)	G34W	186
GC33(L)	G34Y	187
GC33(L)	G34R	188

[0018] Also the invention features an antibody having an activity equivalent to the activity of the antibody described above, wherein one or more amino acid residues are substituted, deleted or added and/or inserted from the amino acid sequences described above.

[0019] Preferably, the antibody of the invention is a humanized antibody.

[0020] Thus, in another aspect, the invention provides a humanized antibody capable of binding to glypican 3.

[0021] In further aspect, the invention provides an antibody capable of binding to a peptide consisting of the sequence of the amino acid residues 524 - 563 of glypican 3.

[0022] Preferably, the antibody of the invention is capable of binding to a peptide consisting of the sequence of the amino acid residues 537 - 563 of glypican 3. More preferably, the antibody of the invention does not bind to a peptide consisting of the sequence of the amino acid residues 550 - 563 of glypican 3.

[0023] Preferably, the antibody is capable of binding to a peptide consisting of the sequence of the amino acid residues 544 - 553 of glypican 3 or a peptide consisting of the sequence of the amino acid residues 546 - 551 of glypican 3.

[0024] In still another aspect, the invention provides an antibody capable of binding to an epitope to which a second antibody is capable of binding, wherein said second antibody comprises a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 143, 144 and 158, respectively. Namely, the antibody of the invention is capable of competing in binding to GPC3 with the second antibody.

[0025] In a preferred embodiment, the antibody of the invention is capable of binding to glypican 3 and has a high CDC activity against a cell expressing glypican 3 and/or has a high ADCC activity against a cell expressing glypican 3.

[0026] In another aspect, the invention provides a polynucleotide coding for a heavy chain variable region or a light chain variable region of the antibody of the invention.

[0027] Preferably, the polynucleotide of the invention has the sequence set forth in SEQ ID NOs: 11-21, 33-43, 55-59, 65-70 and 77-83.

[0028] In still another aspect, the invention provides a cell-growth inhibitor and an anticancer agent comprising as an active ingredient the antibody of the invention. Preferably, the anticancer agent of the invention is used for treatment of hepatoma.

[0029] In further aspect, the invention provides a peptide comprising the sequence of the amino acid residues 524 -

563 of glypican 3, the sequence of the amino acid residues 537 - 563 of glypican 3, the sequence of the amino acid residues 544 - 553 of glypican 3 or the amino acid sequence of the amino acid residues 546 - 551 of glypican 3.

BRIEF DESCRIPTION OF THE DRAWINGS

[0030]

Fig. 1 shows the binding activity of the anti-GPC3 antibody to a CHO cell, a CHO cell expressing full-length GPC3, HepG2 and HuH-7, which was evaluated by flow cytometry. M1E7 (solid line) and M11F1 (dashed line) were used at a concentration of 5 μ g/mL, respectively.

Fig. 2 is a table showing the results of epitope classification by a competitive ELISA. The degrees of competitive inhibition against the binding of the biotinylated anti-GPC3 antibody are indicated by percentage. The epitopes were classified into 5 groups, a to e, according to the competitive inhibition pattern.

Fig. 3 shows the results of evaluating by Western blotting whether an anti-GPC3 antibody binds to the N-terminal fragment of 40 kDa of the soluble form of GPC3 core protein or to the C-terminal fragment of 30 kDa thereof. It was found that L9G11 binds to the N-terminal fragment and M3C11 binds to the C-terminal fragment.

Fig. 4 shows the results of detecting a secreted form of GPC3 is present in the culture supernatant of HepG2 by a sandwich ELISA. It was strongly detected with the combination of antibodies that bind to the N-terminal fragment such as M6B1, M18D4 or M19B11, and it was not strongly detected with an antibody that binds to the C-terminal fragment such as M3C11, M13B3 or M3B8.

Fig. 5 shows the results of immunoprecipitation of the culture supernatant of HepG2 with the use of an anti-GPC3 antibody and detection of a secreted form of GPC3. The medium as a control (lanes 1 and 3) and the culture supernatant of HepG2 (lanes 2 and 4) were immunoprecipitated using M1E7 (lanes 1 and 2) and M10D2 (lanes 3 and 4). Secretory GPC3 was detected by M10D2 that binds to the N-terminal fragment.

Fig. 6 shows the results of analyzing the epitope of the antibodies that bind to the C-terminal fragment of GPC3 by Western blotting with the use of a fusion protein of the C-terminal peptide of GPC3 and GST. The soluble form of GPC3 core protein (lane 1), GST (lane 2), GC-1 (lane 3), GC-2 (lane 4), GC-3 (lane 5), GC-4 (lane 6) and GC-5 (lane 7) were subjected to SDS electrophoresis under reducing conditions, and detected by Western blotting using M3C11 and M11F1.

Fig. 7 shows the results of evaluating the CDC activity of the anti-GPC3 mouse-human chimeric antibody to a CHO cell that expresses GPC3.

Fig. 8 shows the results of evaluating the ADCC activity of the anti-GPC3 mouse-human chimeric antibody to a CHO cell that expresses GPC3 and HepG2.

Fig. 9 shows the results of evaluating the ADCC activity of GC33 to a human hepatoma cell line, HuH-7, using a mouse bone marrow-derived effector cell.

Fig. 10 shows the results of evaluating the antitumor activity of GC33 antibody to a mouse model transplanted with human hepatoma.

Fig. 11 shows the results of evaluating the CDC activity of the mouse-human chimeric antibody GC33 to a CHO cell that expresses GPC3.

Fig. 12 shows the results of evaluating the ADCC activity of the mouse-human chimeric antibody GC33 to HepG2.

Fig. 13 shows GPC3-derived sequences contained in GST-fusion proteins (GC-4, 5, 6, 7, 8, 9, 11, 12, 13 and 14) prepared for analyzing the epitope of GC33.

Fig. 14 shows the results of Western blotting with the use of GC33 after separating GST, GC-7, 8, 9, 11, 12, 13 and 14 by SDS-PAGE under reducing conditions.

Fig. 15 shows the results of evaluating the binding activity of humanized GC33 to GPC3 by an ELISA.

Fig. 16 shows an antibody panel which summarizes isotypes and the results of an ELISA, BIAcore, FACS, an epitope analysis and an immunoprecipitation test for clones derived from a mouse immunized with a soluble form of GPC3.

Fig. 17 shows an antibody panel in which isotypes and the results of an ELISA, FACS and an epitope analysis for clones derived from a mouse immunized with GC-3 are summarized.

Fig. 18 shows the results of evaluating the binding activity of the modified antibodies to the soluble form of GPC3 core protein by an ELISA. Gly34 located at CDR1 in a humanized GC33 L chain variable region was replaced with any of 17 amino acids other than Cys and Met.

Fig. 19 shows the results of evaluating the CDC activity of the mouse-human chimeric antibodies GC33, M3C11, and M1E7 to a CHO cell that expresses full-length GPC3.

Fig. 20 shows the results of evaluating the ADCC activity of the mouse-human chimeric antibodies GC33, M3C11, and M1E7 to a human hepatoma cell line SK-03 that expresses full-length GPC3.

DETAILED DESCRIPTION OF THE INVENTION

Antibody

5 **[0031]** The present invention provides antibodies described in the following (I) to (XI).

(I) An antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: in any of the following (1) to (12):

- 10 (1) SEQ ID NOs: 123, 124 and 125 (GC33),
 (2) SEQ ID NOs: 109, 110 and 111 (M11F1),
 (3) SEQ ID NOs: 106, 107 and 108 (M3B8),
 (4) SEQ ID NOs: 132, 133 and 134 (GC199),
 (5) SEQ ID NOs: 106, 135 and 136 (GC202),
 15 (6) SEQ ID NOs: 126, 127 and 128 (GC179),
 (7) SEQ ID NOs: 129, 130 and 131 (GC194),
 (8) SEQ ID NOs: 103, 104 and 105 (M13B3),
 (9) SEQ ID NOs: 118, 121 and 122 (L9G11),
 (10) SEQ ID NOs: 115, 116 and 117 (M6B1),
 20 (11) SEQ ID NOs: 112, 113 and 114 (M5B9), and
 (12) SEQ ID NOs: 118, 119 and 120 (M10D2).

25 Among the antibodies described in (1) to (12), preferred are the antibodies described in (1) to (8), more preferred are the antibodies described in (1) to (5), and particularly preferred is the antibody described in (1). The antibodies described in (1) to (8) recognize the C-terminal peptide of glypican 3 (a peptide comprising the 374th amino acid to the 580th amino acid of glypican 3); and are useful as a therapeutic antibody. In addition, the antibodies described in (9) to (12) recognize the N-terminal peptide of glypican 3 (a peptide comprising from the 1st amino acid to the 373rd amino acid of glypican 3); and are useful as a diagnostic antibody.

30 (II) An antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: in any of the following (1) to (13):

- 35 (1) SEQ ID NOs: 143, 144 and 158 (GC33),
 (2) SEQ ID NOs: 143, 144 and 145 (M11F1),
 (3) SEQ ID NOs: 140, 141 and 142 (M3B8),
 (4) SEQ ID NOs: 167, 168 and 169 (GC199),
 (5) SEQ ID NOs: 170, 144 and 171 (GC202),
 (6) SEQ ID NOs: 159, 160 and 161 (GC179),
 (7) SEQ ID NOs: 162, 147 and 163 (GC194 (1)),
 40 (8) SEQ ID NOs: 164, 165 and 166 (GC194 (2)),
 (9) SEQ ID NOs: 137, 138 and 139 (M13B3),
 (10) SEQ ID NOs: 155, 156 and 157 (L9G11),
 (11) SEQ ID NOs: 149, 150 and 151 (M6B1),
 (12) SEQ ID NOs: 146, 147 and 148 (M5B9), and
 45 (13) SEQ ID NOs: 152, 153 and 154 (M10D2).

50 Among the antibodies described in (1) to (13), preferred are the antibodies described in (1) to (8), more preferred are the antibodies described in (1) to (5), and particularly preferred is the antibody described in (1). The antibodies described in (1) to (8) recognize the C-terminal peptide of glypican 3 (a peptide comprising from the 374th amino acid to the 580th amino acid of glypican 3); and are useful as a therapeutic antibody. In addition, the antibodies described in (9) to (13) recognize the N-terminal peptide of glypican 3 (a peptide comprising from the 1st amino acid to the 373rd amino acid of glypican 3); and are useful as a diagnostic antibody.

55 (III) An antibody selected from the group consisting of the antibodies described in the following (1) to (13):

- (1) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 123, 124 and 125, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 143, 144 and 158 (GC33),

- (2) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 109, 110 and 111, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 143, 144 and 145 (M11F1),
- (3) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 106, 107 and 108, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 140, 141 and 142 (M3B8),
- (4) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 132, 133 and 134, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 167, 168 and 169 (GC199),
- (5) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 106, 135 and 136, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 170, 144 and 171 (GC202),
- (6) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 126, 127 and 128, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 159, 160 and 161 (GC179),
- (7) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 129, 130 and 131, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 162, 147 and 163 (GC194 (1)),
- (8) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 129, 130 and 131, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 164, 165 and 166 (GC194 (2)),
- (9) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 103, 104 and 105, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 137, 138 and 139 (M13B3),
- (10) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 118, 121 and 122, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 155, 156 and 157 (L9G11),
- (11) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 115, 116 and 117, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 149, 150 and 151 (M6B1),
- (12) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 112, 113 and 114, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 146, 147 and 148 (M5B9),
- (13) an antibody containing heavy chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 118, 119 and 120, and light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 152, 153 and 154 (M10D2).

Among the antibodies described in (1) to (13), preferred are the antibodies described in (1) to (8), more preferred are the antibodies described in (1) to (5), and particularly preferred is the antibody described in (1). The antibodies described in (1) to (8) recognize the C-terminal peptide of glypican 3 (a peptide comprising from the 374th amino acid to the 580th amino acid of glypican 3); and are useful as a therapeutic antibody. In addition, the antibodies described in (9) to (13) recognize the N-terminal peptide of glypican 3 (a peptide comprising from the 1st amino acid to the 373rd amino acid of glypican 3); and are useful as a diagnostic antibody.

(IV) An antibody having a heavy chain variable region described in any of the following (1) to (7):

- (1) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 84 (GC33 VH ver.a),
- (2) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 85 (GC33 VH ver.c),
- (3) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 86 (GC33 VH ver.f),
- (4) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 87 (GC33 VH ver.h),
- (5) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 88 (GC33 VH ver.i),
- (6) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 89 (GC33 VH ver.j), and
- (7) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 90 (GC33 VH ver.k).

Among the antibodies described in (1) to (7), particularly preferred are the antibodies described in (2) to (7).

(V) An antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO:

92 (GC33 VL ver.a).

(VI) An antibody selected from the group consisting of the antibodies described in the following (1) to (7):

- (1) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 84 (GC33 VH ver. a) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a),
- (2) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 85 (GC33 VH ver. c) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a),
- (3) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 86 (GC33 VH ver.f) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a),
- (4) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 87 (GC33 VH ver.h) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a),
- (5) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 88 (GC33 VH ver.i) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a),
- (6) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 89 (GC33 VH ver.j) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a), and
- (7) an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 90 (GC33 VH ver.k) and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 92 (GC33 VL ver.a).

Among the antibodies described in (1) to (7), particularly preferred are the antibodies described in (2) to (7).

(VII) An antibody described in any of the following (1) to (15):

- (1) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 174, 144 and 158,
- (2) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 175, 144 and 158,
- (3) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 176, 144 and 158,
- (4) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 177, 144 and 158,
- (5) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 178, 144 and 158,
- (6) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 179, 144 and 158,
- (7) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 180, 144 and 158,
- (8) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 181, 144 and 158,
- (9) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 182, 144 and 158,
- (10) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 183, 144 and 158,
- (11) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 184, 144 and 158,
- (12) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 185, 144 and 158,
- (13) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 186, 144 and 158,
- (14) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid sequences set forth in SEQ ID NOs: 187, 144 and 158, and
- (15) an antibody containing light chain variable regions having CDRs 1, 2 and 3 consisting of the amino acid

(VIII) An antibody described in any of the following (1) to (15):

- Among the antibodies described in (1) to (15), preferred is the antibody described in (15).

(IX) An antibody described in any of the following (1) to (15):

- 16

- (2) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 192,
- (3) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 193,
- 5 (4) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 194,
- (5) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 195,
- 10 (6) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 196,
- (7) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 197,
- (8) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 198,
- 15 (9) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 199,
- (10) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 200,
- 20 (11) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 201,
- (12) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 202,
- (13) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 203,
- 25 (14) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 204, and
- (15) an antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 205.

30 Among the antibodies described in (1) to (15), preferred is the antibody described in (15).

(X) An antibody having a light chain variable region selected from the group consisting of the light chain variable regions described in the following (1) to (15):

- 35 (1) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 191,
- (2) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 192,
- (3) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 193,
- (4) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 194,
- 40 (5) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 195,
- (6) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 196,
- (7) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 197,
- (8) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 198,
- (9) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 199,
- 45 (10) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 200,
- (11) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 201,
- (12) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 202,
- (13) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 203,
- (14) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 204, and
- 50 (15) a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 205, and a heavy chain variable region selected from the group consisting of the heavy chain variable regions described in the following (1) to (7):

- (1) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 84,
- 55 (2) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 85,
- (3) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 86,
- (4) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 87,
- (5) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 88,
- (6) a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 89, and

Among the antibodies described above, preferred is the antibody having a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 205 and a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 90.

(XI) An antibody, in which one or more amino acids have been replaced, deleted, added and/or inserted in the amino acid sequence described in any one of the above-mentioned (I) to (X), and which has an activity equivalent to that of the antibody described in any of (I) to (X).

[0032] In the present invention, the activity equivalent to that of the antibody described in any of (I) to (X) means that the binding activity to a human glypican 3 antibody or the cytotoxicity activity on a cell that expresses human glypican 3 (e.g., HepG2 or a recombinant CHO cells expressing human glypican 3, etc.) is equivalent.

Humanized antibody

[0033] One preferred embodiment of the antibody according to the present invention is a humanized antibody that binds to glypican 3. The humanized antibody can be prepared by using a known method.

[0034] The humanized antibody is also referred to as a reshaped human antibody, which is made by transplanting the complementarity determining region (CDR) of an antibody of a non-human mammal, for example a mouse antibody, into the CDR of a human antibody. The general recombinant DNA technology for preparation of such antibodies is also known (see European Patent Application EP 125023 and International Patent Application WO 96/02576).

[0035] Specifically, for example, in the case where a CDR is derived from a mouse antibody, a DNA sequence which has been designed to link the CDRs of the mouse antibody with the framework region (FR) of a human antibody is synthesized by the PCR method using several oligonucleotides as primers, which have been prepared so as to have portions overlapping with one another at both ends of the CDR and the FR (see the method described in International Patent Application WO 98/13388).

[0036] As for the framework region of a human antibody to be linked with the CDR, the one which allows a complementarity determining region to form a favorable antigen-binding site is selected. If necessary, an amino acid in the framework region of a variable region of the antibody may be replaced so that the complementarity determining region of a reshaped human antibody may form an appropriate antigen-binding site (Sato, K. et al., Cancer Res. (1993) 53, 851-856).

[0037] The C region of a human antibody may be used as the C region of a chimeric antibody or a humanized antibody, for example, C γ 1, C γ 2, C γ 3, and C γ 4 may be used in the H chain, and C κ and C λ may be used in the L chain. The C region of a human antibody may also be modified in order to improve the stability of the antibody or the production thereof. The human antibody to be used in the humanization may be any isotype of human antibody, for example, IgG, IgM, IgA, IgE and IgD, preferably, IgG, more preferably IgG1 or IgG3, and particularly preferably IgG1. the present invention IgG1 is effective when an antibody is used as an anticancer agent in terms of having a high cytotoxicity activity (Chemical immunology, 65: 88 (1997)).

[0038] In addition, after the humanized antibody is prepared, an amino acid in a variable region (e.g., FR) or a constant region may be replaced with another amino acid.

[0039] The origin of the CDR in a humanized antibody is not particularly limited, and the CDR may be derived from any animals. For example, it is possible to use a sequence derived from a mouse antibody, a rat antibody, a rabbit antibody, a camel antibody or the like. Preferred is a CDR sequence of a mouse antibody.

[0040] With regard to the humanization of an antibody, it is generally difficult to humanize it while maintaining the agonist activity of the original antibody. In the present invention, however, a humanized antibody having an agonist activity equivalent to that of the original mouse antibody was successfully acquired. Since the antigenicity of the humanized antibody in the human body is reduced, it is useful in administering it into the human for a therapeutic purpose.

[0041] Preferred examples of the humanized anti- glypican 3 antibody in the present invention include, for example, an antibody having a heavy chain variable region set forth in SEQ ID NO: 84 (GC33 VH ver.a), SEQ ID NO: 85 (GC33 VH ver.c), SEQ ID NO: 86 (GC33 VH ver.f), SEQ ID NO: 87 (GC33 VH ver.h), SEQ ID NO: 88 (GC33 VH ver.i), SEQ ID NO: 89 (GC33 VH ver.j) or SEQ ID NO: 90 (GC33 VH ver.k) or an antibody having a light chain variable region set forth in SEQ ID NO: 92 (GC33 VL ver. a). Particularly preferred examples thereof include an antibody having a heavy chain variable region set forth in SEQ ID NO: 84 (GC33 VH ver.a), SEQ ID NO: 85 (GC33 VH ver.c), SEQ ID NO: 86 (GC33 VH ver.f), SEQ ID NO: 87 (GC33 VH ver.h), SEQ ID NO: 88 (GC33 VH ver.i), SEQ ID NO: 89 (GC33 VH ver.j) or SEQ ID NO: 90 (GC33 VH ver.k) and a light chain variable region set forth in SEQ ID NO: 92 (GC33 VL ver.a).

[0042] In addition, a preferred example of the humanized anti-glypican 3 antibody includes an antibody having a heavy chain variable region containing the amino acid sequence set forth in SEQ ID NO: 90 and a light chain variable region containing the amino acid sequence set forth in SEQ ID NO: 205.

[0043] A preferred embodiment of the antibody according to the present invention is an antibody that binds to the

epitope to which the antibody set forth in any of the following (1) to (8) binds:

- (1) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 62 and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 73 (GC33),
- (2) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 26 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 48 (M11F1),
- (3) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 25 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 47 (M3B8),
- (4) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 60 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 71 (GC199),
- (5) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 61 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 72 (GC202),
- (6) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 63 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 74 (GC179),
- (7) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 64 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 75 (GC194 (1)), and
- (8) an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 64 and a light chain variable region the amino acid sequence set forth in SEQ ID NO: 76 (GC194 (2)). More preferred is an antibody that binds to the epitope to which the antibody described in any of (1) to (5) binds, and particularly preferred is an antibody that binds to the epitope to which the antibody described in (1) binds.

[0044] The antibody that binds to the epitope to which any of the above-mentioned antibodies bind is useful because it has a particularly high cytotoxicity.

[0045] The antibody described in any of (1) to (7) binds to a region from the 524th amino acid to the 580th amino acid of human glypican 3. In particular, it binds to a region from the 524th amino acid to the 563rd amino acid. The antibody described in any of (1) to (5) binds to a region from the 537th amino acid to the 563rd amino acid of human glypican 3. The antibody described in (1) binds to a region from the 544th amino acid to the 553rd amino acid of human glypican 3. In particular, it binds to a region from the 546th amino acid to the 551st amino acid.

[0046] The antibodies recognizing the above-mentioned epitopes have a high cytotoxicity, therefore they are useful in the treatment of a disease such as cancer. In particular, the antibody which binds to a region from the 546th amino acid to the 551st amino acid is useful as it has a particularly high cytotoxicity

[0047] Accordingly, the present invention includes the antibodies which binds to an epitope in a region from the 524th amino acid to the 580th amino acid of human glypican 3, preferably a region from the 524th amino acid to the 563rd amino acid, more preferably a region from the 537th amino acid to the 563rd amino acid, further more preferably a region from the 544th amino acid to the 553rd amino acid, particularly preferably a region from the 546th amino acid to the 551st amino acid.

[0048] Another preferred embodiment according to the present invention is an antibody that recognizes a region from the 524th amino acid to the 563rd amino acid of human glypican 3 and does not recognize a region from the 537th amino acid to the 563rd amino acid.

[0049] A further preferred embodiment according to the present invention is an antibody that recognizes a region from the 537th amino acid to the 563rd amino acid of human glypican 3 and does not recognize a region from the 550th amino acid to the 563rd amino acid.

[0050] The analysis of an epitope recognized by an antibody can be carried out by a method known to those skilled in the art, for example, by Western blotting described in Examples below.

[0051] The antibody that recognizes the above-mentioned regions as an epitope can be obtained by a method known to those skilled in the art. For example, it can be obtained by preparing a peptide containing an amino acid sequence of a target region based on an amino acid sequence of human glypican 3 and preparing an antibody with the use of the peptide as an immunogene, or by preparing an antibody by a usual method and determining an epitope that the obtained antibody recognizes, and then selecting an antibody that recognizes the target epitope.

[0052] A preferred example of the anti-glypican 3 antibody of the present invention is an antibody having a high ADCC activity or an antibody having a high CDC activity to a cell that expresses glypican 3.

[0053] The phrase "a high ADCC activity" or "a high CDC activity" as used herein means that the antibody of the invention has a higher ADCC activity or a higher CDC activity than that of a known anti-glypican 3 antibody. Known glypican 3 antibodies include, for example, M3C11 and M1E07 described in International Patent Application WO 2004/22739.

[0054] The ADCC activity or the CDC activity can be measured by a method known to those skilled in the art. For example, it can be measured by the chromium release test. Specific conditions of the chromium release test for measuring the ADCC activity are not particularly limited, however, for example, it can be measured using the conditions described

in the Examples below.

[0055] Examples of the cells that express glypican 3 include, for example, a hepatoma cell line such as HepG2, a CHO cell line having a gene encoding glypican 3 incorporated therein and the like. To measure the ADCC activity, it is preferred to use a HepG2 cell line, and to measure the CDC activity, it is preferred to use a recombinant CHO cell line that expresses GPC3. The recombinant CHO cell line that expresses GPC3 may be prepared by any method, however, it can be prepared by, for example, the method described in the Examples below.

[0056] In the case where the anti-glypican 3 antibody is used as an anticancer agent, it is preferred that it has an ADCC activity at the same level as that of an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 62 and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 73 (GC33). In the case where the anti-glypican 3 antibody is used as an anticancer agent, it is preferred that it has a CDC activity at the same level as that of an antibody containing a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 62 and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 73 (GC33).

[0057] Further, the present invention includes an antibody having a high binding activity to glypican 3.

[0058] In the present invention, the binding activity of the antibody to glypican 3 can be measured by using a method known to those skilled in the art. For example, it can be measured by utilizing the surface plasmon resonance with BIAcore. Specifically, a glypican 3 protein is immobilized on a sensor chip to react with an antibody, and the interaction between the antibody and glypican 3 can be calculated as a reaction rate constant from the measurement value. In addition, with regard to the evaluation of the binding activity, an enzyme linked immunosorbent assay (ELISA), an enzyme immunoassay (EIA), a radioimmunoassay (RIA) or a fluorescent antibody technique can be used. For example, in the case where an enzyme immunoassay is used, a sample containing an antibody to be tested, for example, a culture supernatant of a cell producing an antibody to be tested or a purified antibody is added to a plate which has been coated with an antigen to which the antibody to be tested binds. Then, a secondary antibody labeled with an enzyme such as alkaline phosphatase is added, and the plate is incubated and washed. Then, an enzyme substrate such as p-nitrophenyl phosphate is added and the absorbance is measured, whereby an antigen binding activity can be evaluated. The upper limit of the binding activity is not particularly limited. However, for example, the upper limit can be defined within the range which is technically possible by those skilled in the art. It will be appreciated that the range which is technically possible will be expanded by the advancement of technology.

[0059] Further, in the present invention, an amino acid to be deamidated or an amino acid adjacent to an amino acid to be deamidated may be replaced with another amino acid for the purpose of, for example, suppressing deamidation to increase the stability of the antibody. The amino acid to be deamidated includes, asparagine and glutamine, preferably asparagine. An amino acid adjacent to asparagine is not particularly limited and may be any amino acid. It is known that an asparagine-glycine sequence is particularly susceptible to deamidation, thus, glycine is preferred as the amino acid adjacent to asparagine. An amino acid used for replacement is not particularly limited and may be any amino acid other than asparagine and glutamine. Preferred is an amino acid other than valine and proline. Therefore, in the present invention, in the case where the antibody is deamidated, it is preferred to replace the amino acid with an amino acid other than asparagine, glutamine, valine and proline. Suppression of deamidation by amino acid replacement can be carried out with reference to, for example, International Patent Application WO 03/057881. In the case where amino acid replacement is carried out for the purpose of suppression of deamidation, it is preferred that the antigen binding activity before replacement is maintained.

[0060] Another embodiment of stabilization of the antibody includes replacement of glutamic acid with another amino acid. In addition, in the present invention, it was found that, in the case where the 6th amino acid of the heavy chain of an antibody is glutamic acid, the antibody can be significantly stabilized by replacing the glutamic acid with glutamine. Accordingly, the present invention also relates to a method of stabilizing an antibody by replacing the glutamic acid at the 6th position of the heavy chain of the antibody with glutamine. The amino acid numbering of the antibody is known to those skilled in the art (e.g., Kabat, E. A. et al., "Sequences of Proteins of Immunological Interest", US Dept. Health and Human Services 1983).

[0061] The antibody of the invention may be a conjugated antibody in which the antibody is conjugated with various molecules, such as polyethyleneglycol (PEG), radioactive materials and toxin. Such a conjugated antibody may be prepared by chemically modifying the antibody obtained as above. Methods for modifying antibodies have already been established in the art. The antibody of the invention encompasses such a conjugated antibody.

[0062] The antibody of the invention may also be a bispecific antibody (see, for example, Journal of Immunology, 1994, 152, 5368-5374). The bispecific antibody may recognize glypican 3 and another antigen, or may recognize different epitopes on a GPC3 molecule.

[0063] Further, the antibody of the invention may carry a certain protein fused to the N- or C-terminus of the antibody (Clinical Cancer Research, 2004, 10, 1274-1281). The protein to be fused to the antibody may be conveniently selected by those skilled in the art.

[0064] In addition, the antibody of the invention includes an antibody with an enhanced cytotoxicity. Examples of the

antibody with an enhanced cytotoxicity include an antibody lacking fucose, an antibody having bisecting N-acetyl glucosamine (GlcNAc) attached to its sugar chain, and an antibody having altered binding activity for Fc γ receptor obtained by substituting one or more amino acids in the Fc region. Such antibodies with an enhanced cytotoxicity can be prepared by a method known in the art.

Method of preparing antibody

[0065] The antibody that binds to glypican 3 can be prepared by a method known to those skilled in the art. For example, a monoclonal antibody-producing hybridoma can be prepared as follows basically using a known technique. That is, the hybridoma can be prepared by immunizing a mammal in accordance with a usual immunization method using a glypican 3 protein or a cell that expresses glypican 3 as a sensitizing antigen. The thus obtained immunocyte is fused with a known parent cell by a usual cell fusion method, and then selecting a monoclonal antibody-producing cell by a usual screening method.

[0066] Specifically, a monoclonal antibody can be prepared as follows. First, a glypican 3 protein is obtained based on the glypican 3 gene/amino acid sequence shown in SEQ ID NOs: 3 and 4, which is used as a sensitizing antigen to obtain an antibody. More specifically, the gene sequence encoding glypican 3 is inserted into a known expression vector system, and an appropriate host cell is transformed with the vector, and then a target human glypican 3 protein is purified by a known method from the host cell or the culture supernatant.

[0067] Subsequently, this purified glypican 3 protein is used as a sensitizing antigen. Alternatively, a partial peptide of glypican 3 can be used as a sensitizing antigen. In this case, the partial peptide can also be obtained by chemical synthesis according to the amino acid sequence of human glypican 3.

[0068] The epitope on a glypican 3 molecule which is recognized by the anti-glypican 3 antibody of the present invention is not limited to a particular epitope. The anti-glypican 3 antibody may recognize any epitope, as long as the epitope is present on a glypican 3 molecule. Accordingly, any fragment can also be used as an antigen for producing the anti-glypican 3 antibody of the present invention, as long as it contains an epitope that is present on a glypican 3 molecule.

[0069] A mammal to be immunized with a sensitizing antigen is not particularly limited, but it is preferably selected in view of compatibility with a parent cell to be used for cell fusion. For example, rodents such as mice, rats and hamsters, rabbits or monkeys are generally used.

[0070] Immunization of an animal with a sensitizing antigen is carried out in accordance with a known method. For example, immunization is carried out by a general method in which a mammal is injected intraperitoneally or subcutaneously with a sensitizing antigen. Specifically, a sensitizing antigen is diluted with or suspended in an appropriate amount of PBS (Phosphate-Buffered Saline), physiological saline or the like, an appropriate amount of a standard adjuvant such as a Freund's complete adjuvant is mixed with the product if necessary, and then the solution is emulsified and is administered to a mammal several times every 4 to 21 days. In addition, an appropriate carrier can also be used upon immunization with a sensitizing antigen.

[0071] A mammal is immunized as described above, and then an increased level of a target antibody in the serum is confirmed. Subsequently, immunocytes are collected from the mammal, and then subjected to cell fusion. A particularly preferred immunocyte is a splenocyte.

[0072] As a parent partner cell to be fused with the above-mentioned immunocyte, a mammalian myeloma cell is used. Examples of a cell line of the myeloma cell that is preferably used herein include various known cell lines such as P3 (P3x63 Ag8.653) (J. Immunol. (1979) 123, 1548-1550), P3x63 Ag8U.1 (Current Topics in Microbiology and Immunology (1978) 81, 1-7), NS-1 (Kohler, G. and Milstein, C. Eur. J. Immunol. (1976) 6, 511-519), MPC-11 (Margulies, D. H. et al., Cell (1976) 8, 405-415), SP2/0 (Shulman, M. et al., Nature (1978) 276, 269-270), FO (de St. Groth, S. F. et al., J. Immunol. Methods (1980) 35, 1-21), S194 (Trowbridge, I. S. J. Exp. Med. (1978) 148, 313-323) and R210 (Galfre, G. et al., Nature (1979) 277, 131-133).

[0073] Cell fusion of the above-mentioned immunocytes with myeloma cells can be basically carried out in accordance with a known method, for example, the method of Kohler and Milstein et al. (Kohler, G. and Milstein, C., Methods Enzymol. (1981) 73, 3-46).

[0074] More specifically, the above-mentioned cell fusion is carried out in a normal nutrition culture solution in the presence of, for example, a cell-fusion accelerator. As the cell-fusion accelerator, for example, polyethylene glycol (PEG), a hemagglutinating virus of Japan (HVJ) is used. If desired, an adjuvant such as dimethylsulfoxide can be added to further enhance the fusion efficiency.

[0075] The ratio of immunocytes to myeloma cells may be appropriately selected. For example, it is preferred that the number of immunocytes is 1 to 10 times greater than that of myeloma cells. The culture solution to be used for the above-mentioned cell fusion include, for example, a RPMI1640 culture solution or a MEM culture solution which is suitable for the growth of the above-mentioned myeloma cell line, or another normal culture solution which is used for this type of cell culture. Moreover, a serum supplement such as fetal calf serum (FCS) can be used in combination therewith.

[0076] Cell fusion is carried out as follows. Predetermined amounts of the above-mentioned immunocytes and myeloma

cells are mixed well in the above-mentioned culture solution, a PEG (e.g., with an average molecular weight of approximately 1000 to 6000) solution (a general concentration of 30 to 60% (w/v)), which had been pre-heated at approximately 37°C, is added, and then the solution is mixed, whereby a target fusion cell (hybridoma) is formed. Subsequently, an appropriate culture solution is added successively, and then a step of removing the supernatant by centrifugation is repeated to remove a reagent for cell fusion or the like that is unfavorable for the growth of the hybridoma.

[0077] The thus obtained hybridoma is selected by culturing the hybridoma in a standard selective culture solution such as a HAT culture solution (a culture solution containing hypoxanthine, aminopterin and thymidine). Cultivation in the above-mentioned HAT culture solution is continued for a time period sufficient for the cells (unfused cells) other than the target hybridoma to die (normally, several days to several weeks). Subsequently, a standard limiting dilution method is conducted to screen for and monoclonal of hybridoma that produces a target antibody.

[0078] In addition to the method of immunizing a non-human animal with an antigen to obtain hybridoma, a desired human antibody having a binding activity to glypican 3 can also be obtained by sensitizing a human lymphocyte with glypican 3 in vitro, and allowing the sensitized lymphocyte to fuse with a human-derived myeloma cell having a permanent division potential (see JP-B-1-59878). In another method, glypican 3 antigen is administered to a transgenic animal having all the repertoires of human antibody genes to obtain anti-glypican 3 antibody-producing cells, which are then immortalized, and a human antibody for glypican 3 may be obtained from the immortalized anti-glypican 3 antibody-producing cells (see International Patent Applications WO 94/25585, WO 93/12227, WO 92/03918 and WO 94/02602).

[0079] The thus prepared hybridoma that produce a monoclonal antibody can be passage-cultured in a standard culture solution, or can be stored for a long period in liquid nitrogen.

[0080] One example of a method employed to obtain a monoclonal antibody from the hybridoma involves culturing the hybridoma and obtaining a monoclonal antibody from the culture supernatant in accordance with a standard method. Another method involves administering the hybridoma to a mammal that is compatible with the hybridoma to allow it to proliferate, and obtaining a monoclonal antibody from the ascites. The former method is suitable to obtain an antibody of high purity, while the latter method is suitable for the mass production of antibodies.

[0081] It is also possible to prepare a recombinant antibody by cloning the antibody gene from the hybridoma, incorporating the gene into an appropriate vector, introducing the vector into a host, and then allowing the host to produce the recombinant antibody by a genetic engineering technique (e.g., see Vandamme, A. M. et al., Eur. J. Biochem. (1990) 192, 767-775, 1990).

[0082] Specifically, mRNA encoding the variable (V) region of an anti-glypican 3 antibody is isolated from a hybridoma producing the anti-glypican 3 antibody. mRNA is isolated by a known method such as a guanidine ultracentrifugal method (Chirgwin, J. M. et al., Biochemistry (1979) 18, 5294-5299) or an AGPC method (Chomczynski, P. et al., Anal. Biochem. (1987) 162, 156-159), and total RNA is prepared, and then target mRNA is prepared using an mRNA Purification Kit (Pharmacia) or the like. In addition, mRNA can also be directly prepared using a QuickPrep mRNA Purification Kit (Pharmacia).

[0083] The cDNA of the antibody V region is synthesized using a reverse transcriptase from the thus obtained mRNA. cDNA may be synthesized using an AMV Reverse Transcriptase First-strand cDNA Synthesis Kit (SEIKAGAKU CORPORATION) or the like. In addition, for example, a 5'-Ampli FINDER RACE Kit (Clontech), the 5'-RACE method using PCR (Frohman, M. A. et al., Proc. Natl. Acad. Sci. USA (1988) 85, 8998-9002, Belyavsky, A. et al., Nucleic Acids Res. (1989) 17, 2919-2932) can be employed for synthesizing and amplifying cDNA.

[0084] A target DNA fragment is purified from the thus obtained PCR product, and then the vector is introduced into E. coli or the like, and a colony is selected, thereby preparing a desired recombinant vector. The nucleotide sequence of the target DNA is then determined by a known method such as a dideoxynucleotide chain termination method.

[0085] After DNA encoding the V region of the target anti-glypican 3 antibody is obtained, this DNA is incorporated into an expression vector containing DNA encoding the constant region (C region) of the target antibody.

[0086] To produce the anti-glypican 3 antibody used in the present invention, the antibody gene is incorporated into an expression vector so that the gene is expressed under the regulation of the gene expression control region including, for example, an enhancer and a promoter. Next, a host cell is transformed with the expression vector, thereby allowing the host to express the antibody.

[0087] An antibody gene can be expressed by incorporating a polynucleotide encoding the H chain or a polynucleotide encoding the L chain separately into an expression vector, and then simultaneously transforming a host cell with the vectors, or by incorporating polynucleotides encoding the H chain and the L chain into a single expression vector, and then transforming a host cell with the vector (see International Patent Application WO 94/11523).

Polynucleotide

[0088] In another aspect, the present invention provides a polynucleotide encoding a heavy chain variable region or a light chain variable region of the antibody of the present invention. Preferably, the polynucleotide of the present invention

has a nucleotide sequence described in any of SEQ ID NOs: 11-21, 33-43, 55-59, 65-70 and 77-83. In addition, a polynucleotide that is hybridized to the above-mentioned polynucleotide under stringent conditions and encodes an antibody having an activity equivalent to that of the antibody of the present invention is also within the scope of the present invention.

[0089] The polynucleotide of the present invention is not particularly limited as long as it encodes the antibody of the present invention. It is a polymer composed of a plurality of nucleotides, such as deoxyribonucleic acids (DNA) or ribonucleic acids (RNA). It may contain a base other than a naturally occurring base. The polynucleotide of the present invention can be used for producing an antibody by a genetic engineering technique. In addition, the polynucleotide of the present invention can be used as a probe to screen for an antibody having a function equivalent to that of the antibody of the present invention. That is, a polynucleotide encoding the antibody of the present invention or a partial fragment thereof may be used as a probe to obtain DNA that is hybridized to the polynucleotide under stringent conditions and encodes an antibody having an activity equivalent to that of the antibody of the present invention by techniques such as a hybridization technique, a gene amplification technique (e.g., PCR). Such DNA is also included in the polynucleotide of the present invention.

[0090] The hybridization technique (Sambrook, J. et al., *Molecular Cloning* 2nd ed., 9.47-9.58, Cold Spring Harbor Lab. Press, 1989) is well known to those skilled in the art. Examples of the hybridization conditions include, for example, low stringent conditions. The low stringent conditions are, for example, the conditions of 42°C, 0.1x SSC and 0.1% SDS, preferably the conditions of 50°C, 0.1x SSC and 0.1% SDS when washing is performed after hybridization. More preferred examples of the hybridization conditions include, for example, high stringent conditions. The high stringent conditions are, for example, the conditions of 65°C, 5x SSC and 0.1% SDS. Under these conditions, it can be expected that a polynucleotide having a higher homology can be efficiently obtained under higher temperature. Incidentally, there are plural factors that affect the stringency of hybridization, such as temperature and the concentration of salt, and those skilled in the art can achieve a similar stringency by appropriately selecting these factors.

[0091] An antibody functionally equivalent to the antibody of the present invention encoded by a polynucleotide obtained by such a hybridization technique and a gene amplification technique usually has a high homology with the antibody in terms of the amino acid sequence. The antibody of the present invention also includes an antibody that is functionally equivalent to the antibody of the present invention and has a high homology with the amino acid sequence of the antibody. A high homology means generally at least 50% or higher identity, preferably 75% or higher identity, more preferably 85% or higher identity, and further more preferably 95% or higher identity at the amino acid level. To determine the homology of polypeptides, the algorithm described in the literature (Wilbur, W. J. and Lipman, D. J., *Proc. Natl. Acad. Sci. USA* (1983) 80, 726-730) may be employed.

[0092] The present invention also provides a vector containing the polynucleotide of the present invention. Such a vector can be used for preparing the antibody of the present invention. As for the vector of the present invention, in the case where *E. coli* is used as a host, for example, it is not particularly limited as long as it has "ori" for use in amplification in *E. coli* to produce and amplify the vector in a large amount in *E. coli* (e.g., JM109, DH5 α , HB101 or XL1Blue), and has a marker gene for selecting a transformed *E. coli* (e.g., a drug resistance gene that can be identified by a drug such as ampicillin, tetracycline, kanamycin or chloramphenicol). Examples of the vector include M13-series vectors, pUC-series vectors, pBR322, pBluescript, pCR-Script and the like. In addition, pGEM-T, pDIRECT, and pT7 can also be used for subcloning and extracting cDNA as well as the vectors described above.

[0093] As the vector of the present invention, an expression vector is particularly useful. For example, an expression vector to be expressed in *E. coli* should have the above characteristics to be amplified in *E. coli*. In addition, in the case where *E. coli*, such as JM109, DH5 α , HB101, or XL1-Blue is used as a host cell, it is indispensable that the vector should have a promoter, for example, lacZ promoter (Ward et al., *Nature* (1989) 341, 544-546; *FASEB J.* (1992) 6, 2422-2427), araB promoter (Better et al., *Science* (1988) 240, 1041-1043), T7 promoter or the like, that can efficiently express the desired product in *E. coli*. Examples of such a vector include pGEX-5X-1 (Pharmacia), "QIAexpress system" (Qiagen), pEGFP, pET (in this case, the host is preferably BL21 which expresses T7 RNA polymerase) and the like, as well as the vectors described above.

[0094] In addition, the vector may also contain a signal sequence for polypeptide secretion. As for the signal sequence for protein secretion, in the case where a polypeptide is produced in the periplasm of *E. coli*, the pelB signal sequence (Lei S. P. et al., *J. Bacteriol* (1987) 169, 4379) can be used. Introduction of the vector into a host cell can be carried out by using, for example, the calcium chloride method and the electroporation method.

[0095] In addition to *E. coli*, for example, expression vectors derived from mammals (e.g., pcDNA3 (Invitrogen) and pEGF-BOS (Nucleic Acids Res. (1990) 18(17), p5322), pEF and pCDM8), expression vectors derived from insect cells (e.g., "Bac-to-BAC baculovirus expression system" (GIBCO BRL) and pBacPAK8), expression vectors derived from plants (e.g., pMH1 and pMH2), expression vectors derived from animal viruses (e.g., pHSV, pMV and pAdexLow), expression vectors derived from retroviruses (e.g., pZIPneo), expression vectors derived from yeast (e.g., "Pichia Expression Kit" (Invitrogen), pNV11 and SP-Q01), and expression vectors derived from *Bacillus subtilis* (e.g., pPL608 and pKTH50) can be used as the vector of the present invention.

[0096] For the purpose of expressing the vector in an animal cell such as a CHO cell, a COS cell, an NIH3T3 cell or the like, it is indispensable for the vector to have a promoter required for expression in a cell such as SV40 promoter (Mulligan et al., Nature (1979) 277, 108), MMTV-LTR promoter, EF1 α promoter (Mizushima et al., Nucleic Acids Res. (1990) 18, 5322), CMV promoter or the like, and more preferably to have a marker gene (such as a drug resistance gene that can be identified by a drug such as neomycin or G418) for selecting transformation into the cell. Examples of the vector having such characteristics include, for example, pMAM, pDR2, pBK-RSV, pBK-CMV, pOPRSV and pOP13.

[0097] Further, for the purpose of stably expressing a gene and, at the same time, amplifying the gene copy numbers in the cell, a vector (e.g., pCHO1, etc.) having the DHFR gene is introduced into the CHO cell deficient in the nucleic acid synthetic pathway to complement the deficiency and is amplified with methotrexate (MTX). In addition, for the purpose of transient expression of a gene, transformation is effected with a vector (such as pcD) having the origin of replication for SV40 using a COS cell having on the chromosome a gene that expresses the SV40 T antigen. As the origin of replication, the one derived from a polyoma virus, an adenovirus, a bovine papilloma virus (BPV) and the like can also be used. Further, for the amplification of gene copy numbers in the host cell system, the expression vector can include, as a selectable marker, the aminoglycoside transferase (APH) gene, the thymidine kinase (TK) gene, E. coli xanthine guaninephosphoribosyl transferase (Ecogpt) gene, the dihydrofolate reductase (dhfr) gene and the like.

[0098] To prepare the antibody of the present invention, the vector is introduced into a host cell. The host cell into which the vector is introduced is not particularly limited, but includes, for example, E. coli or any of various animal cells. For example, the host cell can be used as a production system for the production or expression of the antibody of the present invention. As for the production system of polypeptide preparation, there are an in vitro production system and an in vivo production system. In vitro production system include a production system which employs eukaryotic cells and a production system which employs prokaryotic cells.

[0099] In the case where the eukaryotic cell is used, for example, an animal cell, a plant cell or a fungal cell can be used. Known animal cells include a mammalian cell such as a CHO cell (J. Exp. Med. (1995) 108, 945), a COS cell, a 3T3 cell, a myeloma cell, a baby hamster kidney (BHK) cell, a HeLa cell and a Vero cell, an amphibian cell such as a Xenopus oocyte (Valle, et al., Nature (1981) 291, 358-340), or an insect cell such as Sf9, Sf21, and Tn5. In the present invention, CHO-DG44, CHO-DXB11, a COS7 cell, a BHK cell are preferably used. Among the animal cells, for the purpose of performing a large amount of expression, a CHO cell is particularly preferred. Introduction of the vector into the host cell can be carried out by, for example, the calcium phosphate method, the DEAE-dextran method, the cationic ribosome DOTAP (Boehringer Mannheim), the electroporation method, the lipofection method or the like.

[0100] As for the plant cell, for example, a cell derived from Nicotiana tabacum is known as a protein production system, which may be subjected to callus culture. Examples of the fungal cells include yeast such as the genus Saccharomyces, more specifically Saccharomyces cerevistae and Saccharomyces pombe, and filamentous fungi such as the genus Aspergillus, more specifically Aspergillus niger.

[0101] In the case where the prokaryotic cell is used, production system using a bacterial cell may be employed. Examples of the bacterial cells include Escherichia coli (E. coli) such as JM109, DH5 α and HB101, and Bacillus subtilis.

Preparation of recombinant antibody

[0102] The antibody of the present invention can be prepared by culturing the above-mentioned host cells. The antibody can be obtained by culturing in vitro a cell transformed with a desired polynucleotide. Cultivation can be carried out in accordance with a known method. Culture media for animal cells include, for example, DMEM, MEM, RPMI 1640, and IMDM. A serum supplement such as FBS or fetal calf serum (FCS) may be used in combination, or serum-free medium can be used. The pH during the cultivation is preferably about 6 to 8. Cultivation is usually carried out at about 30 to 40°C for about 15 to 200 hours with, as needed, medium change, aeration, and agitation.

[0103] On the other hand, systems for producing a polypeptide in vivo include, for example, a production system which employs an animal and a production system which employs a plant. The target polynucleotide is introduced into such an animal or a plant, and the polypeptide is produced in the body of the animal or the plant and recovered. The term "host cell" as used herein encompasses such an animal and a plant.

[0104] When the animal is used, production systems employing a mammal or an insect are available. As the mammal, goats, pigs, sheep, mice and cattle can be used (Vicki Glaser, SPECTRUM Biotechnology Applications, 1993). A transgenic animal can also be used as a mammal.

[0105] For example, the target polynucleotide is prepared as a fusion gene with a gene encoding a polypeptide which is inherently produced in the milk such as goat β casein. Then, the DNA fragment containing this fusion gene is injected into a goat embryo, and the embryo is transplanted into a female goat. The target antibody can be obtained from the milk produced by the transgenic goat borne to the goat which received the embryo or the offspring thereof. To increase the amount of milk containing the antibody produced by the transgenic goat, hormone may be given to the transgenic goat as needed. (Ebert, K. M. et al., Bio/Technology (1994) 12, 699-702).

[0106] In addition, as an insect, for example, a silkworm can be used. In the case where a silkworm is used, a silkworm

is infected with a baculovirus into which the polynucleotide encoding the target antibody has been inserted. The target antibody can be obtained from the body fluid of the silkworm (Susumu, M. et al., *Nature* (1985) 315, 592-594).

[0107] In the case where a plant is used, for example, tobacco can be used. In the case where tobacco is used, a polynucleotide encoding the target antibody is inserted into an expression vector for a plant, for example pMON 530, and then the vector is introduced into a bacterium such as *Agrobacterium tumefaciens*. Then, tobacco such as *Nicotiana tabacum* is infected with the bacterium, whereby the target antibody can be obtained from the leaves of the tobacco (Julian, K. -C. Ma et al., *Eur. J. Immunol.* (1994) 24, 131-138).

[0108] The thus obtained antibody can be isolated from the inside or the outside (culture medium, etc.) of the host cell and then can be purified to a substantially pure and uniform antibody. Separation and purification of the antibody may be carried out by a separation and a purification method usually used in purification of polypeptides. For example, polypeptides can be separated and purified by any methods including chromatography columns, filtration, ultrafiltration, salting-out, solvent precipitation, solvent extraction, distillation, immunoprecipitation, SDS-polyacrylamide gel electrophoresis, isoelectric focusing, dialysis, recrystallization, and a combination thereof.

[0109] Examples of the chromatography include, for example, affinity chromatography, ion exchange chromatography, hydrophobic chromatography, gel-filtration, reverse phase chromatography, adsorption chromatography (Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Ed Daniel R. Marshak et al., Cold Spring Harbor Laboratory Press, 1996). These chromatographies can be carried out using a liquid phase chromatography such as HPLC and FPLC. Examples of a column to be used for affinity chromatography include a protein A column or a protein G column. One example of the protein A column is Hyper D, POROS, Sepharose F. F. (Pharmacia).

[0110] Further, before or after purification of the antibody, the antibody can be modified or peptides can be partially removed as needed by allowing a suitable protein-modifying enzyme to act on. The protein-modifying enzyme for this purpose include, for example, trypsin, chymotrypsin, lysyl endopeptidase, protein kinase, glucosidase.

Diagnostic method

[0111] In another aspect, the present invention provides a method of diagnosing a disease such as cancer by detecting GPC3 protein in a test sample with the use of the antibody of the present invention.

[0112] The detection used herein includes quantitative detection and non-quantitative detection. The non-quantitative detection include, for example, determination of merely whether or not GPC3 protein is present, determination of whether or not a specific amount or more of GPC3 protein is present, determination for comparison of the amount of GPC3 protein with that of another sample (e.g., a control sample). The quantitative detection includes determination of the concentration of GPC3 protein, determination of the amount of GPC3 protein.

[0113] The test sample is not particularly limited as long as it is a sample that may possible contain GPC3 protein, however, preferred is a sample collected from the body of a living organism such as a mammal, and more preferred is a sample collected from human. Specific examples of the test sample may include, for example, blood, interstitial fluid, plasma, extravascular fluid, cerebral fluid, joint fluid, pleural fluid, serum, lymph fluid, saliva, preferably blood, serum and plasma. In addition, a sample obtained from the test sample such as culture solution of cells collected from the body of the living organism is also included in the test sample of the present invention.

[0114] The cancer to be diagnosed is not particularly limited, and specific examples may include liver cancer, pancreatic cancer, lung cancer, colon cancer, mammary cancer, prostate cancer, leukemia and lymphoma, preferably liver cancer. GPC3 to be detected is not particularly limited, and may be either full-length GPC3 or a fragment thereof. In the case where a fragment of GPC3 is detected, it may be either the N-terminal fragment or the C-terminal fragment, however, the N-terminal fragment is preferred. In addition, the GPC3 protein may also be a heparan sulfate-added GPC3 or a GPC3 core protein.

[0115] The method of detecting GPC3 protein contained in a test sample is not particularly limited, however, detection is preferably performed by an immunological method with the use of an anti-GPC3 antibody. Examples of the immunological method include, for example, a radioimmunoassay, an enzyme immunoassay, a fluorescence immunoassay, a luminescence immunoassay, immunoprecipitation, a turbidimetric immunoassay. Preferred is an enzyme immunoassay, and particularly preferred is an enzyme-linked immunosorbent assay (ELISA) (e.g., a sandwich ELISA). The above-mentioned immunological method such as an ELISA can be carried out by a method known to those skilled in the art.

[0116] A general detection method with the use of an anti-GPC3 antibody comprises immobilizing an anti-GPC3 antibody on a support, adding a test sample thereto, incubating the support to allow the anti-GPC3 antibody and GPC3 protein to bind to each other, washing the support, and detecting the GPC3 protein binding to the support via the anti-GPC3 antibody to detect GPC3 protein in a test sample.

[0117] The binding between the anti-GPC3 antibody and the GPC3 protein is generally carried out in a buffer. Buffers used in the invention include, for example, a phosphate buffer, a Tris buffer. Incubation is carried out under the conditions generally employed, for example, at 4 °C to room temperature for 1 hour to 24 hours. The washing after incubation can be carried out by any method as long as it does not inhibit the binding between the GPC3 protein and the anti-GPC3

antibody, using for example a buffer containing a surfactant such as Tween 20.

[0118] In the method of detecting GPC3 protein of the present invention, a control sample may be provided in addition to a test sample to be tested for GPC3 protein. The control samples include a negative control sample that does not contain GPC3 protein and a positive control sample that contains GPC3 protein. In this case, it is possible to detect GPC3 protein in the test sample by comparing the result obtained with the negative control sample that does not contain GPC3 protein with the result obtained with the positive control sample that contains GPC3 protein. It is also possible to quantitatively detect GPC3 protein contained in the test sample by obtaining the detection results of the control samples and the test sample as numerical values, and comparing these numerical values.

[0119] One preferred embodiment of detecting GPC3 protein binding to the support via an anti-GPC3 antibody is a method using an anti-GPC3 antibody labeled with a detectable label. For example, GPC3 protein may be detected by contacting the test sample with an anti-GPC3 antibody immobilized on the support, washing the support, and then detecting GPC3 with the use of the labeled antibody that specifically binds to GPC3 protein.

[0120] The labeling of an anti-GPC3 antibody can be carried out by a generally known method. Examples of the detectable label known to those skilled in the art include a fluorescent dye, an enzyme, a coenzyme, a chemiluminescent substance or a radioactive substance. Specific examples may include radioisotopes (^{32}P , ^{14}C , ^{125}I , ^3H , ^{131}I and the like), fluorescein, rhodamine, dansyl chloride, umbelliferone, luciferase, peroxidase, alkaline phosphatase, β -galactosidase, β -glucosidase, horseradish peroxidase, glucoamylase, lysozyme, saccharide oxidase, microperoxidase, biotin and the like. In the case where biotin is used as a detectable label, it is preferred that a biotin-labeled antibody is added, and then avidin conjugated to an enzyme such as alkaline phosphatase is further added.

[0121] Specifically, a solution containing an anti-GPC3 antibody is added to a support such as a plate to allow the anti-GPC3 antibody to be immobilized. After washing, the plate is blocked with, for example, BSA in order to prevent the nonspecific binding of a protein. The plate is washed again, and then the test sample is added to the plate. After being incubated, the plate is washed, and then the labeled anti-GPC3 antibody is added. After being incubated appropriately, the plate is washed, and then the labeled anti-GPC3 antibody remaining on the plate is detected. The detection of the protein can be carried out by a method known to those skilled in the art. For example, in the case where the antibody is labeled with a radioactive substance, the protein may be detected by liquid scintillation or the RIA method. In the case where the antibody is labeled with an enzyme, the protein may be detected by adding a substrate and detecting an enzymatic change of the substrate such as color development with an absorbance reader. In the case where the antibody is labeled with a fluorescent substance, the protein may be detected with the use of a fluorometer.

[0122] A particularly preferred embodiment of the method of detecting GPC3 protein of the present invention is a method using an anti-GPC3 antibody labeled with biotin and avidin. Specifically, a solution containing an anti-GPC3 antibody is added to a support such as a plate to allow the anti-GPC3 antibody to be immobilized thereon. After washing, the plate is blocked with, for example, BSA in order to prevent the nonspecific binding of a protein. The plate is washed again, and then the test sample is added to the plate. After being incubated, the plate is washed, and then the biotin-labeled anti-GPC3 antibody is added. After being incubated appropriately, the plate is washed, and then avidin conjugated to an enzyme such as alkaline phosphatase or peroxidase is added. After being incubated, the plate is washed, and then a substrate of the enzyme conjugated to avidin is added. Then, GPC3 protein is detected by means of the enzymatic change of the substrate as an indicator.

[0123] Another embodiment of the method of detecting GPC3 protein of the present invention is a method using a primary antibody that specifically binds to GPC3 protein and a secondary antibody that specifically binds to the primary antibody. For example, the test sample is brought into contact with an anti-GPC3 antibody immobilized on the support, the support is incubated and washed, and the bound GPC3 protein after washing is detected with a primary anti-GPC3 antibody and a secondary antibody that specifically binds to the primary antibody. In this case, the secondary antibody is preferably labeled with a detectable label.

[0124] Specifically, a solution containing an anti-GPC3 antibody is added to a support such as a plate to allow the anti-GPC3 antibody to be immobilized thereon. After washing, the plate is blocked with, for example, BSA in order to prevent the nonspecific binding of a protein. The plate is washed again, and then the test sample is added to the plate. After being incubated, the plate is washed, and then a primary anti-GPC3 antibody is added. After being incubated appropriately, the plate is washed, and then a secondary antibody that specifically binds to the primary antibody is added. After being incubated appropriately, the plate is washed, and then the secondary antibody remaining on the plate is detected. The detection of the secondary antibody can be carried out by the above-mentioned method.

Pharmaceutical composition

[0125] In another aspect, the present invention provides a pharmaceutical composition containing the antibody of the present invention. The pharmaceutical composition containing the antibody of the present invention is useful in the treatment and/or prevention of a disease associated with cell proliferation such as cancer, and particularly it is useful in the treatment and/or prevention of liver cancer. In the case where the antibody of the present invention is used as a

pharmaceutical composition, the antibody can be formulated into a dosage form by a method known to those skilled in the art. For example, the pharmaceutical composition can be used parenterally in the form of an injection of a sterile solution or a suspension with water or another pharmaceutically acceptable solution. For example, the antibody can be formulated into a dosage form by appropriately mixing it with a pharmaceutically acceptable carrier or solvent, such as sterile water, physiological saline, a plant-oil, an emulsifier, a suspension, a surfactant, a stabilizer, a flavor, an excipient, a vehicle, a preservative, a binder to prepare a unit dosage form required for generally accepted Drug Implementation. The amount of active ingredients in these preparations is selected to allow for administration of a suitable dosage within the indicated range.

[0126] A sterile composition for injection can be formulated by using a vehicle such as distilled water for injection in accordance with the general Drug Implementation.

[0127] Examples of the aqueous solution for injection include, for example, physiological saline, glucose, and other isotonic liquids including adjuvants, such as D-sorbitol, D-mannose, D-mannitol and sodium chloride. They can be used in combination with a suitable solubilizer, such as an alcohol, specifically ethanol, a polyalcohol such as propylene glycol and polyethylene glycol, and a non-ionic surfactant such as Polysorbate 80 (TM) and HCO-50.

[0128] Sesame oil or soybean oil can be used as a oleaginous liquid and may be used in combination with benzyl benzoate or benzyl alcohol as a solubilizer. It may be formulated with a buffer such as a phosphate buffer or a sodium acetate buffer, a pain-killer such as procaine hydrochloride, a stabilizer such as benzyl alcohol or phenol, or an antioxidant. The prepared injection is generally filled into a suitable ampule.

[0129] The method of administration is preferably parenteral, and specific examples thereof include injection, transnasal administration, transpulmonary administration, transdermal administration and the like. The injection formulation may be administered systemically or topically by intravenous injection, intramuscular injection, intraperitoneal injection, subcutaneous injection or the like.

[0130] The method of administration can be appropriately selected according to the age and the symptoms of a patient. For example, one dose of the pharmaceutical composition containing the antibody or the polynucleotide encoding the antibody can be selected from the range of 0.0001 mg to 1,000 mg per kg of body weight. Alternatively, for example, the dose can be selected from the range of 0.001 mg to 100,000 mg/body per patient, although it is not always limited to these numerical values. The dose and the method of administration vary according to the body weight, the age and the symptoms of a patient, and are appropriately selected by those skilled in the art.

[0131] All patents and references cited in this specification are incorporated by reference. All the contents disclosed in the specifications and drawings of Japanese Patent Application No. 2004-203637, on which the application claims priority, are incorporated herein by reference.

EXAMPLE

[0132] The present invention will be described in more detail with reference to Examples below. However, the present invention is not limited to these Examples.

Example 1

cDNA cloning of human glypican 3 (GPC3)

[0133] A full-length cDNA encoding human GPC3 was amplified by PCR reaction with an Advantage 2 kit (CLONTECH) using 1st stranded cDNA prepared by a usual method from a colon cancer cell line, Caco2, as a template. More specifically, 50 μ L of a reaction mixture containing 2 μ L of cDNA derived from Caco2, 1 μ L of a sense primer (GATATC-ATGGCCG-GGACCGTGCGCACC GCGT: SEQ ID NO: 1), 1 μ L of an antisense primer (GCTAGC-TCAGTGCACCAGGAAGAA-GAAGCAC: SEQ ID NO: 2), 5 μ L of Advantage 2 10x PCR buffer, 8 μ L of dNTP mix (1.25 mM) and 1.0 μ L of Advantage polymerase Mix was subjected to 35 cycles consisting of 94 °C for 1 minute, 63 °C for 30 seconds and 68 °C for 3 minutes. The amplified product from the PCR reaction was inserted into a TA vector, pGEM-T Easy, using pGEM-T Easy Vector System I (Promega). The sequence was confirmed by using an ABI 3100 DNA sequencer. In this way, a cDNA encoding full-length human GPC3 was isolated. The sequence shown in SEQ ID NO: 3 indicates the nucleotide sequence of human GPC3 gene and the sequence shown in SEQ ID NO: 4 indicates the amino acid sequence of human GPC3 protein.

Example 2

Preparation of soluble form of GPC3

[0134] As immunoprotein for the generation of an anti-GPC3 antibody, a soluble form of GPC protein was prepared, in which a hydrophobic region at the C-terminal side (564-580 amino acids) was deleted.

[0135] By using the full-length human GPC3 cDNA as a template, a PCR reaction was carried out using an antisense primer (ATA GAA TTC CAC CAT GGC CGG GAC CGT GCG C: SEQ ID NO: 5) and a sense primer, to which an EcoRI recognition sequence and a Kozak sequence were added, (ATA GGA TCC CTT CAG CGG GGA ATG AAC GTT C: SEQ ID NO: 6). The obtained PCR fragment (1711 bp) was cloned into pCXND2-Flag. The pCXND2-Flag was designed to express a Flag-tagged protein by inserting the region for DHFR gene expression of pCHO1 (Hirata et al., FEBS letter 1994; 356; 244-248) into the HindIII site of pCXN2 (Niwa et al., Gene 1991; 108; 193-199) and adding a Flag tag sequence to the downstream of the multicloning site. The constructed expression plasmid DNA was introduced into a CHO cell line, DXB11, and a CHO cell line highly expressing the soluble form of GPC3 was obtained by selection with 500 µg/mL Geneticin. The large-scale cultivation of the CHO cell line highly expressing the soluble form of GPC3 was carried out using a 1700-cm² roller bottle, and the culture supernatant was recovered for the antibody purification. The culture supernatant was applied to a DEAE sepharose Fast Flow column (Amersham) and, after washing, the antibody was eluted with a buffer containing 500 mM NaCl, and affinity purified using Anti-Flag M2 agarose affinity gel (SIGMA). The elution was carried out with 200 µg/mL FLAG peptide. After the eluate was concentrated with Centriprep-10 (Millipore), FLAG peptide was removed by gel filtration using Superdex 200 HR 10/30 (Amersham). Lastly, the filtrate was concentrated using a DEAE sepharose Fast Flow column and eluted with PBS (containing 500 mM NaCl) without Tween 20 to effect buffer exchange.

Example 3

Preparation of soluble form of GPC3 core protein

[0136] GPC3 is modified by heparan sulfate to become a macromolecule. To eliminate an antibody against heparan sulfate in a screening for an anti-GPC3 antibody, a soluble form of GPC3 core protein that had a point mutation in the heparan sulfate-binding site was prepared and used in the screening.

[0137] Using the above-mentioned soluble form of GPC3 (1-563) as a template, a cDNA in which Ser residues at the 495th and 509th positions were replaced with Ala was prepared by the assembly PCR method, in which primers were designed to add His tag to the C-terminus. The obtained cDNA was cloned into pCXND3 vector. The pCXND3 was constructed by inserting the DHFR gene expressing region of pCHO1 in the HindIII site of pCXN2. The constructed expression plasmid DNA was introduced into DXB11 cell line and a CHO cell line highly expressing a soluble form of GPC3 core protein was obtained by selection with 500 µg/mL Geneticin.

[0138] The large-scale cultivation was carried out using a 1700-cm² roller bottle and the culture supernatant was recovered for antibody purification. The culture supernatant was applied to a Q sepharose Fast Flow column (Amersham). After washing, the antibody was eluted with a phosphate buffer containing 500 mM NaCl, and affinity purified using a Chelating sepharose Fast Flow column (Amersham). The antibody was eluted with a gradient of 10 to 150 mM imidazole. Lastly, the eluate was concentrated using a Q sepharose Fast Flow column and, eluted with a phosphate buffer containing 500 mM NaCl.

[0139] SDS polyacrylamide gel electrophoresis under reducing conditions showed three bands of 70 kDa, 40 kDa and 30 kDa. The result of amino acid sequencing using an ABI492 protein sequencer (Applied Biosystems) indicated that the 30 kDa band corresponded to the amino acid sequence of the 359th and its downstream or the 375th and its downstream of GPC3, suggesting that GPC3 was cleaved between Arg358 and Ser359 or between Lys374 and Val375, hence, it was separated into 40 kDa of the N-terminal fragment and 30 kDa of the C-terminal fragment.

Example 4

Preparation of CHO cell line expressing full-length human GPC3

[0140] To obtain a cell line for evaluating a binding activity using flow cytometry, a CHO cell line expressing full-length GPC3 was established.

[0141] Ten microgram of a full-length human GPC3 gene expression vector and 60 µL of SuperFect (QIAGEN) were mixed. After a complex was formed, gene introduction was carried out by adding it to a CHO cell line, DXB11. After a 24-hour cultivation in a CO₂ incubator, selection was started using αMEM (GIBCO BRL) containing Geneticin at a final concentration of 0.5 mg/mL and 10% FBS. The resulting Geneticin-resistant colonies were collected and cell cloning was carried out by the limiting dilution method. Each cell clone was solubilized and the expression of full-length human GPC3 was confirmed by Western blotting using an anti-GPC3 antibody. In this way, a stably expressing cell line was obtained.

Example 5

Evaluation of binding activity by ELISA

[0142] The soluble form of GPC3 core protein was diluted to 1 µg/mL with a coating buffer (0.1 mol/L NaHCO₃ (pH 9.6), 0.02% (w/v) NaN₃) and added to an immunoplate and left at 4 °C overnight to coat the plate. After the plate was blocked with a dilution buffer (50 mmol/L Tris-HCl (pH 8.1), 1 mmol/L MgCl₂, 150 mmol/L NaCl, 0.05% (v/v) Tween 20, 0.02% (w/v) NaN₃, 1% (w/v) BSA), an anti-GPC3 antibody was added and left at room temperature for 1 hour. After washing with a rinse buffer (0.05% (v/v) Tween 20, PBS), an anti-mouse IgG antibody (ZYMED) labeled with alkaline phosphatase was added and left at room temperature for 1 hour. After washing with the rinse buffer, SIGMA 104 (SIGMA) diluted to 1 mg/mL with a substrate buffer (50 mmol/L NaHCO₃ (pH 9.8), 10 mmol/L MgCl₂) was added and left at room temperature for 1 hour for color development. Then the absorbance (at 405 nm, reference wavelength of 655 nm) was measured using a Benchmark Plus (BIO-RAD).

Example 6

Immunization with soluble form of GPC3 and selection of hybridoma

[0143] Since human GPC3 and mouse GPC3 show a high homology of 94% at the amino acid level, it was considered difficult to obtain an anti-GPC3 antibody if a normal mouse was immunized. Therefore, an autoimmune disease mouse, MRL/MpJUmCrj-lpr/lpr mouse, (hereinafter referred to as MRL/lpr mouse, purchased from Charles River Japan, Inc.) was used as an immunized animal. Immunization was started at the age of 7 weeks or 8 weeks. For the first immunization, a soluble form of GPC3 was prepared at 100 µg/head and emulsified using Freund's complete adjuvant (FCA, Becton Dickinson) and subcutaneously administered. Two weeks later, a soluble form of GPC3 was prepared at 50 µg/head and emulsified using Freund's incomplete adjuvant (FIA, Becton Dickinson) and subcutaneously administered. After that, an additional immunization was carried out every other week for 5 times in total. To two of the immunized mice, a soluble form of GPC3 was diluted with PBS to 50 µg/head, and then administered intravenously via the tail as the final immunization. On the fourth day after the final immunization, the spleen was excised to obtain a spleen cell, which was mixed with a mouse myeloma cell, P3-X63Ag8U1 (P3U1, purchased from ATCC), at a ratio of 2:1. Cell fusion was carried out by gradually adding PEG 1500 (Roche Diagnostic). RPMI 1640 medium (GIBCO BRL) was carefully added to dilute PEG 1500, and after PEG 1500 was removed by centrifugation, the cells were suspended in RPMI 1640 medium containing 10% FBS and inoculated into a 96-well culture plate at 100 µL/well. On the next day, RPMI 1640 medium containing 10% FBS, 1x HAT media supplement (SIGMA) and 0.5x BM-Condimed H1 Hybridoma cloning supplement (Roche Diagnostic) (hereinafter referred to as HAT medium) was added at 100 µL/well. After 2, 3 and 5 days, half of the culture solution was replaced with the HAT medium. After 7 days, screening was carried out using the culture supernatant. The screening was carried out by an ELISA using an immunoplate coated with the soluble form of GPC3 core protein. A positive clone was monocloned by the limiting dilution method. As a result, 11 clones of antibodies (M3C11, M13B3, M1E7, M3B8, M11F1, L9G11, M19B11, M6B1, M18D4, M5B9 and M10D2) that have a strong binding activity against GPC3 were obtained.

Example 7

Isotype determination and purification of anti-GPC3 antibody

[0144] Isotype was determined by an antigen-dependent ELISA using an Immunopure Monoclonal Antibody Isotyping Kit I (PIERCE). The purification of antibodies was carried out as follows. The culture supernatant of hybridoma cultured with the HAT medium supplemented with FBS (Ultra low IgG) (GIBCO BRL) was adsorbed to Hi Trap ProteinG HP (Amersham), and washed with a binding buffer (20 mM sodium phosphate (pH 7.0)). The antibody was eluted with an elution buffer (0.1 M glycine-HCl (pH 2.7)). The eluate was immediately neutralized with a neutralization buffer (1 M Tris-HCl (pH 9.0)), and dialyzed against PBS for day and night for buffer exchange.

Example 8

Evaluation of binding activity by ELISA

[0145] In order to conveniently evaluate the binding activity of the anti-GPC3 antibody thus obtained, concentration-dependent binding of the antibody was detected against an immunoplate containing the soluble form of GPC3 core protein immobilized thereon. A 3-fold dilution series (12 dilutions in total) of the purified antibody at a concentration of

10 $\mu\text{g/mL}$ was added, and an anti-mouse IgG antibody was added as the secondary antibody. Color development was carried out using SIGMA 104. Since the degree of color development varies depending on the color development time, data measured precisely after 1 hour was analyzed. Every antibody showed a concentration-dependent color development. The correlation between the concentration of antibody and the degree of color development was plotted and an approximate curve was obtained by using an analyzing software, GraphPad Prism. Its EC50 value was determined as the index of the binding activity. EC50 values for all clones are shown in Fig. 16.

Example 9

Evaluation of binding activity by flow cytometry

[0146] Cells were dissociated with 1 mM EDTA pH 8.0 (GIBCO)/PBS and suspended in FACS buffer (1% FBS/PBS) at 1×10^6 cells/mL. The suspension was dispensed to a Multiscreen-HV Filter Plate (Millipore) at 100 μL /well and the supernatant was removed by centrifugation. An anti-GPC3 antibody diluted to an appropriate concentration was added and reacted on ice for 30 minutes. The cells were washed once with FACS buffer and an FITC-labeled anti-mouse IgG antibody was added and reacted on ice for 30 minutes. After the reaction, the cells were centrifuged at 500 rpm for 1 minute, and the supernatant was removed. The cells were suspended in 400 μL of FACS buffer and subjected to flow cytometry. EPICS ELITE ESP (Beckman Coulter) was used as a flow cytometer. A gate was set on the living cell population with the histogram of forward scatter and side scatter. As shown in Fig. 1, an anti-GPC3 antibody (M3C11, M11F1) bound strongly to the CHO cell expressing GPC3 and did not bind to the parent CHO cell, indicating that the antibody specifically binds to GPC3 presented on the cell membrane. In addition, the antibody showed the binding activity to a hepatoma cell line, HepG2 (purchased from ATCC) and HuH-7 (purchased from Health Science Research Resources Bank), suggesting that the antibody may specifically recognize hepatoma. The binding activity of the clones derived from the mouse immunized with a soluble form of GPC3 measured by flow cytometry is shown in Fig. 16, where the X-mode values of histogram at the concentration of antibody of 5 $\mu\text{g/mL}$ are indicated.

Example 10

Epitope classification by competitive ELISA

[0147] The obtained antibodies were classified according to the epitopes by a competitive ELISA. The antibodies were biotinylated using a Biotin Labeling Kit (Roche). The soluble form of GPC3 core protein was diluted to 1 $\mu\text{g/mL}$ with the coating buffer and added to a plate at 100 μL /well and stored at 4°C overnight to coat the plate. On the next day, 200 μL of the substrate buffer was added for blocking. The plate was left at 4°C overnight or longer and an anti-GPC3 antibody was added to the plate at 100 μL /well and reacted at room temperature for 1 hour. After that, without washing of the plate, 10 μL of 10 $\mu\text{g/mL}$ biotin-labeled anti-GPC3 antibody was added and further reacted for 1 hour. The plate was washed with 300 μL /well of the rinse buffer for 3 times. AP-streptavidin conjugate (ZYMED) was diluted to 1000-fold with the dilution buffer and added at 100 μL /well and reacted at room temperature for 1 hour. The plate was washed with 300 μL /well of the rinse buffer for 5 times. SIGMA 104 was diluted to 1 mg/mL with the substrate buffer and added at 100 μL /well. After incubating for 1 hour at room temperature, the absorbance (at 405 nm, reference wavelength of 655 nm) was measured.

[0148] The results of the competitive ELISA are shown in Fig. 2. As for the antibody that competitively inhibited the binding of the biotinylated antibody by 50% or more, it was considered that its epitopes are located close together in the three-dimensional conformation. As a result of classification according to the competitive inhibition pattern of color development against the binding of the 8 types of biotinylated antibodies, the 11 clones derived from the mouse immunized with a soluble form of GPC3 were classified into 5 groups (a, b, c, d and e) (Fig. 16).

Example 11

Epitope classification by Western blotting

[0149] The soluble form of GPC3 core protein was applied to a 10% SDS-PAGE mini (TEFCO) and electrophoresed under reducing conditions. It was transferred to Immobilon-P (Millipore) using Trans-Blot SD Semi-Dry Electrophoretic Transfer Cell (BIO-RAD). After the membrane was briefly washed with TBS-T (0.05% Tween 20, TBS), it was shaken in TBS-T containing 5% skim milk for 1 hour. The membrane was shaken in TBS-T for about 10 minutes, then each anti-GPC3 antibody diluted with TBS-T containing 1% skim milk was added and the membrane was shaken for 1 hour. The membrane was washed with TBS-T and shaken in a solution of HRP-anti-mouse IgG antibody (Amersham) diluted with TBS-T containing 1% skim milk for 1 hour, and then washed with TBS-T. Color development was carried out using ECL-

Plus (Amersham) and detected using Hyperfilm ECL (Amersham).

[0150] As shown in Fig. 3, L9G11 was determined to be an antibody binding to the N-terminal side because it bound to the band of about 40 kDa. M3C11 was determined to be an antibody binding to the C-terminal side because it bound to the band of about 30 kDa. All the antibodies belonging to c, d or e group based on the competitive ELISA bound to the N-terminal side, and all those belonging to a or b groups bound to the C-terminal side (Fig. 16). L9G11 had higher detection sensitivity in Western blotting than the other antibodies that bind to the N-terminal side, suggesting that this antibody is a useful for detecting the N-terminal fragment by Western blotting.

Example 12

Detection of secreted form of GPC3

[0151] Since it was found that GPC3 is cleaved at the 358th amino acid residue or the 374th amino acid residue, the inventors hypothesized that a secreted form of GPC3 is secreted into the blood of a patient with liver cancer. Therefore, a GPC3 sandwich ELISA system was constructed in order to detect a secretory form of GPC3.

[0152] An immunoplate was coated with an anti-GPC3 antibody at 10 $\mu\text{g/mL}$ and blocked by the substrate buffer. After the immunoplate was stored for several hours at room temperature or overnight at 4 °C, the culture supernatant of HepG2 was added and incubated for 1 hour at room temperature. The immunoplate was washed with 300 μL /well of the rinse buffer for 3 times, and a biotin-labeled anti-GPC3 antibody diluted to 10 $\mu\text{g/mL}$ was added and incubated for 1 hour at room temperature. The immunoplate was washed with 300 μL /well of the rinse buffer for 3 times, and AP-streptavidin was added and incubated for 1 hour at room temperature. The immunoplate was washed with 300 μL /well of the rinse buffer for 5 times. Color development was carried out using AMPAK (DAKO) in accordance with the attached protocol and the absorbance was measured using a microplate reader. The antibodies binding to the N-terminal side (M6B1, M19B11 and M18D4) and those binding to the C-terminal side (M3C11, M13B3 and M3B8) were combined to construct five sandwich ELISA systems. Each of these combinations showed an equivalent sensitivity in the standard curve using the secreted form of GPC3. These systems were evaluated using the culture supernatant of HepG2. The secreted form of GPC3 was detected at a high concentration of about 1 $\mu\text{g/mL}$ with a combination of the antibodies binding to the N-terminal side (Fig. 4). The concentration detected with a combination of the antibodies binding to the C-terminal side was low, suggesting that the N-terminal fragment was dominantly present in the secreted form of GPC3.

[0153] Subsequently, the culture supernatant of HepG2 was immunoprecipitated using an anti-GPC3 antibody to detect the secreted form of GPC3. In the case where M10D2 that binds to the N-terminal fragment was used, the secreted form of GPC3 of 40 kDa was detected (Fig. 5). On the other hand, in the case where M1E7 that binds to the C-terminal fragment was used, the secreted form of GPC3 was not detected. The immunoprecipitation test was carried out for all the obtained GPC3 antibodies. Every antibody binding to the N-terminal fragment strongly detected the secreted form of GPC3, while the secreted form of GPC3 was not detected or was weakly detected with the use of the antibodies binding to the C-terminal fragment (Fig. 16). The antibody that can detect the secreted form of GPC3 by immunoprecipitation is expected to be useful as an antibody for diagnosing hepatoma. In addition, the antibody that can hardly detect the secreted form of GPC3 is expected to be useful in the development of a therapeutic antibody having an ADCC activity and a CDC activity, because such an antibody may migrate to the hepatoma lesion without being trapped in the secreted form of GPC3 present in the blood.

Example 13

Cloning of variable region of anti-GPC3 antibody

[0154] A variable region of the anti-GPC3 antibody was amplified by the RT-PCR method using the total RNA extracted from an anti-GPC3 antibody-producing hybridoma. The total RNA was extracted from 1×10^7 cells of the hybridoma with the use of RNeasy Plant Mini Kits (QIAGEN). By using 1 μg of the total RNA, the 5'-terminal gene fragment was amplified with the use of a SMART RACE cDNA Amplification Kit (CLONTECH) and any of the following synthetic oligonucleotides:

a synthetic oligonucleotide MHC-IgG1 complementary to the sequence of a mouse IgG1 constant region:

GGG CCA GTG GAT AGACAG ATG (SEQ ID NO: 7)

;

a synthetic oligonucleotide MHC-IgG2a complementary to the sequence of a mouse IgG2a constant region:

CAG GGG CCA GTG GAT AGA CCG ATG (SEQ ID NO: 8)

;

a synthetic oligonucleotide MHC-IgG2b complementary to the sequence of a mouse IgG2b constant region:

CAG GGG CCA GTG GAT AGA CTG ATG (SEQ ID NO: 9)

; and

a synthetic oligonucleotide kappa complementary to the sequence of a mouse kappa chain constant region:

GCT CAC TGG ATG GTG GGA AGA TG (SEQ ID NO: 10).

[0155] A reverse transcription reaction was carried out at 42°C for 1 hour and 30 minutes. The PCR mixture (50 µL) contained 5 µL of 10x Advantage 2 PCR buffer, 5 µL of 10x Universal Primer A Mix, 0.2 mM dNTPs (dATP, dGTP, dCTP and dTTP), 1 µL of Advantage 2 Polymerase Mix (all from CLONTECH), 2.5 µL of the reverse transcription reaction product and 10 pmol of the synthetic oligonucleotide MHC-IgG1, MHC-IgG2a, MHC-IgG2b or kappa. PCR was carried out with 5 cycles consisting of 94°C for 30 seconds, 94°C for 5 seconds and 72°C for 3 minutes, 5 cycles consisting of 94°C for 5 seconds, 70°C for 10 seconds and 72°C for 3 minutes, and 25 cycles consisting of 94°C for 5 seconds, 68°C for 10 seconds and 72°C for 3 minutes. Lastly, the reaction product was heated at 72°C for 7 minutes. Each PCR product was purified from the agarose gel using a QIAquick Gel Extraction Kit (QIAGEN), cloned into pGEM-T Easy vector (Promega), and the nucleotide sequence was determined.

[0156] The nucleotide sequences of the H chain variable regions of M3C11, M13B3, M1E7, M3B8, M11F1, M19B11, M6B1, M18D4, M5B9, M10D2 and L9G11 are shown in SEQ ID NOs: 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 and 21, respectively, the amino acid sequences thereof are shown in SEQ ID NOs: 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32, respectively. The nucleotide sequences of the L chain variable regions thereof are shown in SEQ ID NOs: 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 and 43, respectively, and the amino acid sequences thereof are shown in SEQ ID NOs: 44, 45, 46, 47, 48, 49, 50, 51, 52, 53 and 54, respectively.

Example 14

Epitope classification using GST-fusion protein

[0157] To carry out a detail analysis of the epitopes for the antibodies binding to the C-terminal fragment, fusion proteins of successively shortened C-terminal peptides of GPC3 with GST, namely GC-1 (from Ser495 to Lys563), GC-2 (from Gly510 to Lys563), GC-3 (from Ala524 to Lys563), GC-4 (from Gly537 to Lys563) and GC-5 (from Ser550 to Lys563) were prepared. The C-terminal region of GPC3 was cloned into pGEX-4T-3 (Amersham) to construct a plasmid DNA in which the C-terminal region of GPC3 is ligated to the C-terminal side of GST. The plasmid DNA was introduced into DH5α, whereby a transformant was obtained. Then, IPTG was added at 1 mM to a culture of the transformant in the logarithmic growth phase to induce the expression of a GST-fusion protein. The bacterial cells were collected after 2 hours cultivation. The cells were homogenized by sonication, and centrifuged at 35,000 rpm for 30 minutes with XL-80 ultracentrifuge (Beckman, 70.1 Ti rotor). Then, the culture supernatant was recovered and purified with GST Purification Modules (Amersham). The thus purified GST-fusion proteins were separated by SDS-PAGE under reducing conditions, and Western blotting was carried out with the anti-GPC3 antibodies (Fig. 6). M3C11 and M1E7 detected GC-1 and GC-2, while they did not detect GC-3, GC-4 and GC-5, indicating that the epitopes of these antibodies are contained in the region of GC-2, and that the region of GC-3 is not sufficient. M3B8 and M11F1 detected GC-1 GC-2, GC-3 and GC-4, while they did not detect GC-5, indicating that the epitopes of these antibodies are contained in the region of GC-4, and that the region of GC-5 is not sufficient. The minimum region of the GST-fusion protein to which each antibody can bind is listed in the column headed "Western blotting" of Fig. 16.

Example 15

Preparation of anti-GPC3 mouse-human chimeric antibody

[0158] The sequences of the H chain and the L chain variable regions of the anti-GPC3 antibodies were ligated to the sequences of a human IgG1 and a kappa chain constant regions. PCR was carried out by using a synthetic oligonucleotide,

which is complementary to the 5'-terminal nucleotide sequence of the H chain variable region of each antibody and has a Kozak sequence, and a synthetic oligonucleotide, which is complementary to the 3'-terminal nucleotide sequence and has a NheI site. The obtained PCR product was cloned into pB-CH vector in which a human IgG1 constant region was inserted into pBluescript KS(+) vector (Toyobo). The mouse H chain variable region and the human H chain (γ 1 chain) constant region are ligated via the NheI site. The prepared H chain gene fragment was cloned into an expression vector, pCXND3. On the other hand, PCR was carried out by using a synthetic oligonucleotide, which is complementary to the 5'-terminal nucleotide sequence of the L chain variable region of each antibody and has a Kozak sequence, and a synthetic oligonucleotide, which is complementary to the 3'-terminal nucleotide sequence and has a BsiWI site. The obtained PCR product was cloned into pB-CL vector in which the human kappa chain constant region was inserted into pBluescript KS(+) vector (Toyobo). The human L chain variable region and the constant region are ligated via the BsiWI site. The prepared L chain gene fragment was cloned into an expression vector, pUCAG. This pUCAG vector was obtained by cloning a 2.6 kbp fragment obtained by digesting pCXN (Niwa et al., Gene 1991; 108: 193-200) with a restriction enzyme BamHI into the BamHI site of pUC19 vector (Toyobo).

[0159] To prepare an expression vector for an anti-GPC3 mouse-human chimeric antibody, a gene fragment was obtained by digesting the pUCAG vector containing the L chain gene fragment with a restriction enzyme HindIII (Takara Shuzo), and cloned into the HindIII site of the pCXND3 containing the H chain gene. This plasmid will express a neomycin-resistance gene, DHFR gene and an anti-GPC3 mouse-human chimeric antibody in an animal cell.

[0160] A CHO cell line (DG44 cell line) stably expressing the antibody was prepared as follows. The gene was introduced into the cells by the electroporation method using Gene Pulser II (Bio-Rad). A mixture obtained by mixing 25 μ g of the expression vector for each anti-GPC3 mouse-human chimeric antibody and 0.75 mL of a solution of CHO cells suspended in PBS (1×10^7 cell/mL) was cooled on ice for 10 minutes, and transferred to a cuvette. Then, a pulse was applied at 1.5 kV and a capacitance of 25 μ FD. After a 10-minute recovery period at room temperature, the electroporated cells were suspended in 40 mL of CHO-S-SFM II medium (Invitrogen) containing 1xHT supplement (Invitrogen). The suspension was diluted to 50-fold with the same medium, and dispensed to a 96-well culture plate at 100 μ L/well. After a 24-hour culture in a CO₂ incubator (5% CO₂), Geneticin (Invitrogen) was added at 0.5 mg/mL and the cells were cultured for 2 weeks. The culture supernatant was taken from the well having a Geneticin resistant transformed cell colony and the amount of IgG was measured by the concentration determination method described below. A high-producing cell line was successively expanded to obtain a cell line that stably expresses an anti-GPC3 mouse-human chimeric antibody. The cell line was cultured at a large-scale and the culture supernatant was collected. The purification of each anti-GPC3 mouse-human chimeric antibody was carried out using Hi Trap ProteinG HP (Amersham).

Example 16

Measurement of complement-dependent cytotoxicity activity (CDC activity)

16.1 Preparation of human albumin veronal buffer (HAVB)

[0161] In milli-Q water, 12.75 g of NaCl (highest grade, Wako Pure Chemicals), 0.5625 g of Na-Barbital (highest grade, Wako Pure Chemicals) and 0.8625 g of Barbital (highest grade, Wako Pure Chemicals) were dissolved to a final volume of 200 mL and autoclaved at 121°C for 20 minutes. Then, 100 mL of autoclaved hot milli-Q water was added. The pH was 7.43 (recommended pH: 7.5). The solution was used as a 5x veronal buffer. In 50 mL of milli-Q water, 0.2205 g of CaCl₂·2H₂O (highest grade, Junsei Chemical) was dissolved to a final concentration of 0.03 mol/L, which was used as a CaCl₂ solution. In 50 mL of milli-Q water, 1.0165 g of MgCl₂·6H₂O (highest grade, Junsei Chemical) was dissolved to a final concentration of 0.1 mol/L, which was used as a MgCl₂ solution. In milli-Q water, 100 mL of the 5x veronal buffer, 4 mL of human serum albumin (25% Buminat (registered trademark), the concentration of human serum albumin: 250 mg/mL, Baxter Healthcare), 2.5 mL of the CaCl₂ solution, 2.5 mL of the MgCl₂ solution, 0.1 g of KCl (highest grade, Junsei Chemical) 0.5 g of glucose (D(+)-glucose, anhydrous glucose, highest grade, Wako Pure Chemicals) were dissolved to a final volume of 500 mL, which was used as HAVB. After filter sterilization, the HAVB was stored at a preset temperature of 5°C.

16.2 Preparation of target cell

[0162] The CHO cell expressing full-length human GPC3 prepared in Example 4 was cultured in α -MEM medium containing nucleic acid (+) (GIBCO) supplemented with 10% FBS and 0.5 mg/mL Geneticin (GIBCO). The cells were dissociated from the dish using a cell dissociation buffer (Invitrogen Corp), dispensed to each well of a 96-well flat-bottomed plate (Falcon) at 1×10^4 cells/well, and cultured for 3 days. After the cultivation, 5.55 MBq of chromium-51 was added, and the cells were cultured in a 5% carbon dioxide gas incubator at 37°C for 1 hour. These cells were washed with HAVB twice, and 50 μ L of HAVB was added and used as a target cell.

16.3 Chromium release test (CDC activity)

[0163] Each chimeric antibody was diluted with HAVB to make a 40 µg/mL antibody solution. To the target cell, 50 µL of each antibody solution was added, and left on ice for 15 minutes. Subsequently, to each well, 100 µL of the human serum from the peripheral blood of a healthy volunteer, which had been diluted with HAVB, was added to a final concentration of 25% (the final concentration of antibody: 10 µg/mL), and left in a 5% carbon dioxide gas incubator at 37°C for 90 minutes. After the plate was centrifuged, 100 µL of the supernatant was collected from each well, the radioactivity was measured using a gamma counter. The specific chromium release rate was obtained by the following formula.

$$\text{Specific chromium release rate (\%)} = (A - C) \times 100 / (B - C)$$

"A" represents the radioactivity (cpm) in each well, "B" represents the mean value of the radioactivities (cpm) in the wells in which 100 µL of 2% NP-40 aqueous solution (Nonidet P-40, Code No. 252-23, Nacalai Tesque) and 50 µL of HAVB were added to the target cell, and "C" represents the mean value of the radioactivities (cpm) in the wells in which 150 µL of HAVB was added to the target cell. The test was carried out in triplicate and the mean value and the standard deviation were calculated for CDC activity (%).

[0164] The results are shown in Fig. 7. Among 9 types of the anti-GPC3 chimeric antibodies, M3B8 and M11F1, which are an antibody recognizing the C-terminal side, showed a strong CDC activity against the CHO cell expressing GPC3, however, the CDC activity was not observed in the other antibodies. M3B8 and M11F1 belong to the group called "b" based on the competitive ELISA, and an important epitope for showing a strong CDC activity could be found.

Example 17

Measurement of ADCC activity using PBMC derived from human peripheral blood17.1 Preparation of human PBMC solution

[0165] The heparinized peripheral blood obtained from a healthy volunteer was diluted to 2-fold with PBS(-), and overlaid on Ficoll-Paque TM PLUS (Amersham). After centrifugation at 500 x g for 30 minutes at 20°C, the middle layer, which is the mononuclear leukocyte fraction, was collected. The cells were washed 3 times, suspended in 10% FBS/RPMI and used as a human PBMC solution.

17.2 Preparation of target cell

[0166] The HepG2 cells cultured in 10% FBS/RPMI 1640 medium were dissociated from the dish using Trypsin-EDTA (Invitrogen), dispensed to each well of a 96-well U-bottomed plate (Falcon) at 1×10^4 cells/well, and cultured for 2 days. The CHO cell expressing full-length human GPC3 prepared in Example 4 was cultured in α -MEM nucleic acids (+) medium (GIBCO) supplemented with 10% FBS and 0.5 mg/mL Geneticin (GIBCO). The cells were dissociated from the dish using a cell dissociation buffer (Invitrogen Corp), dispensed to each well of a 96-well flat-bottomed plate (Falcon) at 1×10^4 cells/well, and cultured for 3 days. Chromium-51 (5.55 MBq) was added to each cell and the cells were cultured in a 5% carbon dioxide gas incubator at 37°C for 1 hour. These cells were washed with the medium once, and 50 µL of 10% FBS/RPMI 1640 medium was added and used as a target cell.

17.3 Chromium release test (ADCC activity)

[0167] To the target cell, 50 µL of an antibody solution prepared at different concentrations was added, and reacted on ice for 15 minutes. Subsequently, 100 µL of the human PBMC solution was added at 5×10^5 cells/well, and cells were cultured in a 5% carbon dioxide gas incubator at 37°C for 4 hours. After the cultivation, the plate was centrifuged, and the radioactivity in 100 µL of the culture supernatant was measured using a gamma counter. The specific chromium release rate was obtained by the following formula.

$$\text{specific chromium release rate (\%)} = (A - C) \times 100 / (B - C)$$

"A" represents the mean value of the radioactivities (cpm) in each well, "B" represents the mean value of the radioactivities

(cpm) in the wells in which 100 μ L of 2% NP-40 aqueous solution (Nonidet P-40, Code No. 252-23, Nacalai Tesque) and 50 μ L of 10% FBS/RPMI medium were added to the target cell, and "C" represents the mean value of the radioactivities (cpm) in the wells in which 150 μ L of 10% FBS/RPMI medium was added to the target cell. The test was carried out in triplicate and the mean value and the standard deviation were calculated for ADCC activity (%). The results are shown in Fig. 8. Among 9 types of the anti-GPC3 chimeric antibodies, the antibodies recognizing the C-terminal side had a tendency of showing a strong ADCC activity.

Example 18

Immunization with GC-3 and selection of hybridoma

[0168] Among the obtained anti-GPC3 antibodies, only M11F1 and M3B8 showed a strong CDC activity, indicating that the CDC activity is epitope dependent. To obtain an antibody having both ADCC activity and CDC activity, a GST-fusion protein containing the epitope for M11F1 and M3B8, referred to as GC-3, was used for immunization. A large amount of GC-3 was purified by the above-mentioned method. The buffer was changed to PBS by gel filtration using Superdex 75 (Amersham). The obtained product was used as immunoprotein. using three Balb/c mice (purchased from Charles River Japan, Inc.) and three MRL/lpr mice were immunized with GC-3 in accordance with the above-mentioned method. For the first immunization, GC-3 was prepared at 100 μ g/head and emulsified using FCA, which was subcutaneously administered. Two weeks later, GC-3 was prepared at 50 μ g/head and emulsified using FIA, which was subcutaneously administered. After the fifth immunization, the final immunization (50 μ g/head) was carried out for all mice by intravenously administering the immunoprotein via the tail. After cell fusion, hybridoma were screened by an ELISA using an immunoplate coated with the soluble form of GPC3 core protein. A positive clone was monocloned by the limiting dilution method. As a result, 5 clones of antibodies (GC199, GC202, GC33, GC179 and GC194) that have a strong binding activity against GPC3 were obtained.

[0169] The antibody was purified from the culture supernatant of the hybridoma using Hi Trap proteinG HP, and analyzed in accordance with the above-mentioned method. The EC50 value was calculated by an ELISA using an immunoplate coated with the soluble form of GPC3 core protein, and the X-mode value of histogram at 5 μ g/mL was measured by flow cytometry (Fig. 17). According to the epitope classification by a competitive ELISA, the antibodies were classified into the group b (GC199, GC202 and GC33) and a new epitope group f (GC179 and GC194). The epitope classification using the GST-fusion proteins indicated that GC199, GC202 and GC33 detected GC-1, GC-2, GC-3 and GC-4, but did not detect GC-5, suggesting that the epitopes for these antibodies are contained in the region of GC-4 in the same manner as the epitopes for M11F1 and M3B8, and that the region of GC-5 is not sufficient. On the other hand, GC179 and GC194 detected GC-1, GC-2 and GC-3, but did not detect GC-4 and GC-5, suggesting that the epitopes for these antibodies are contained in the region of GC-3, and that the region of GC-4 is not sufficient. The minimum region of the GST-fusion protein to which each antibody can bind is listed in the column headed "Western blotting" of Fig. 17.

[0170] The H chain and the L chain variable regions of GC199, GC202, GC33, GC179 and GC194 were cloned in accordance with the above-mentioned method, and their sequences were determined. As for the L chain of GC194, 2 types of sequences were cloned. The nucleotide sequences of the H chain variable regions of GC199, GC202, GC33, GC179 and GC194 are shown in SEQ ID NOs: 55, 56, 57, 58 and 59, respectively, and the amino acid sequences thereof are shown in SEQ ID NOs: 60, 61, 62, 63 and 64, respectively. The nucleotide sequences of the L chain variable regions of GC199, GC202, GC33, GC179, GC194(1) and GC194(2) are shown in SEQ ID NOs: 65, 66, 67, 68, 69 and 70 respectively, and the amino acid sequences thereof are shown in SEQ ID NOs: 71, 72, 73, 74, 75 and 76, respectively.

[0171] Further, these amino acid sequences were examined for homology by comparing with the database of the amino acid sequences of known antibodies, whereby their CDR regions were determined as follows.

Antibody	CDR	Amino Acid Sequence	SEQ ID NO
M13B3(H)	CDR1	NYAMS	103
	CDR2	AINNNGDDTYLDTVKD	104
	CDR3	QGGAY	105
M3B8(H)	CDR1	TYGMGVG	106
	CDR2	NIWWYDAKYNSDLKS	107
	CDR3	MGLAWFAY	108
M11F1(H)	CDR1	IYGMGVG	109

EP 1 674 111 A1

Table continued

Antibody	CDR	Amino Acid Sequence	SEQ ID NO
	CDR2	NIWWNDDKYNSALKS	110
	CDR3	IGYFYFDY	111
M5B9(H)	CDR1	GYWMH	112
	CDR2	AIYPGNSDTNYNQKFKG	113
	CDR3	SGDLTGGLAY	114
M6B1(H)	CDR1	SYAMS	115
	CDR2	AINSNGGTTYYPDTMKD	116
	CDR3	HNGGYENYGWFAY	117
M10D2(H)	CDR1	SYWMH	118
	CDR2	EIDPSDSYTYYNQKFRG	119
	CDR3	SNLGDGHYRFPAPFY	120
L9G11(H)	CDR1	SYWMH	118
	CDR2	TIDPSDSETHYNLQFKD	121
	CDR3	GAFYSSYSYWAWFAY	122
GC33(H)	CDR1	DYEMH	123
	CDR2	ALDPKTGDTAYSQKFKG	124
	CDR3	FYSYTY	125
GC179(H)	CDR1	INAMN	126
	CDR2	RIRSESNNYATYYGDSVKD	127
	CDR3	EVTTSFAY	128
GC194(H)	CDR1	ASAMN	129
	CDR2	RIRSKSNYAIYYADSVKD	130
	CDR3	DPGYYGNPWFAY	131
GC199(H)	CDR1	DYSMH	132
	CDR2	WINTETGEPTYADDFKG	133
	CDR3	LY	134
GC202(H)	CDR1	TYGMGVG	106
	CDR2	NIWWHDDKYNSALKS	135
	CDR3	IAPRYNKYEGFFAF	136
M13B3(L)	CDR1	KSSQSLLSDGKTYLN	137
	CDR2	LVSKLDS	138
	CDR3	WQGTHFPLT	139
M3B8(L)	CDR1	KASQDINNYLS	140
	CDR2	RANRLVD	141
	CDR3	LQCDEFPPWT	142
M11F1(L)	CDR1	RSSQSLVHSNGNTYLH	143
	CDR2	KVSNRFS	144
	CDR3	SQSTHVPWT	145

Table continued

Antibody	CDR	Amino Acid Sequence	SEQ ID NO
M5B9(L)	CDR1	RSSKSLHSHNGITYLY	146
	CDR2	QMSNLAS	147
	CDR3	AQNLELPYT	148
M6B1(L)	CDR1	KASQDINKNII	149
	CDR2	YTSTLQP	150
	CDR3	LQYDNLPR	151
M10D2(L)	CDR1	RASHSISNHL	152
	CDR2	YASQSI	153
	CDR3	QQSNIWSLT	154
L9G11(L)	CDR1	RASESVEYYGTSLMQ	155
	CDR2	GASNVES	156
	CDR3	QSRKVPYT	157
GC33(L)	CDR1	RSSQSLVHSHNGNTYLH	143
	CDR2	KVSNRFS	144
	CDR3	SQNTHPPT	158
GC179(L)	CDR1	KSSKSLHSHNGNTYLN	159
	CDR2	WMSNLAS	160
	CDR3	MQHIEPPT	161
GC194(L)1	CDR1	RSSKSLHSHYDITYLY	162
	CDR2	QMSNLAS	147
	CDR3	AQNLELPPT	163
GC194(L)2	CDR1	SASSSVSYMY	164
	CDR2	DTSNLAS	165
	CDR3	QQWSSYPLT	166
GC199(L)	CDR1	KSSQSLHSDGKTFLN	167
	CDR2	LVSRLDS	168
	CDR3	CQGTHFPRT	169
GC202(L)	CDR1	RSSQSIVHSHNGNTYLE	170
	CDR2	KVSNRFS	144
	CDR3	FQGSHPWT	171

Example 19

Measurement of ADCC activity using mouse bone marrow derived effector cell

19.1 Preparation of mouse bone marrow derived effector cell solution

[0172] Bone marrow cells were collected from the femur of an SCID mouse (CLEA Japan, Inc., male, 10 weeks old), and suspended in 10% FBS/RPMI 1640 medium at 5×10^5 cells/mL. Mouse GM-CSF (PeproTech) and human IL-2 (PeproTech) were added at 10 ng/mL and 50 ng/mL, respectively, and the cells were cultured in a 5% carbon dioxide gas incubator at 37°C for 5 days. After the cultivation, the cells were scraped off with a scraper and washed with the

medium once. Then, the cells were suspended in 10% FBS/RPMI 1640 medium at 5×10^6 cells/mL, and used as a mouse bone marrow derived effector cell solution.

19.2 Preparation of target cell

[0173] A human hepatoma cell line, HuH-7, was maintained and subcultured with DMEM medium (SIGMA) containing 10% FBS (ThermoTrace). The cells were dissociated from the dish using Cell Dissociation Buffer (Invitrogen), dispensed to each well of a 96-well U-bottomed plate (Falcon) at 1×10^4 cells/well, and cultured for 1 day. After the cultivation, 5.55 MBq of chromium-51 was added, and the cells were cultured in a 5% carbon dioxide gas incubator at 37°C for 1 hour. These cells were washed with the medium once, and 50 μ L of 10% FBS/RPMI 1640 medium was added and used as a target cell.

19.3 Chromium release test (ADCC activity)

[0174] To the target cell, 50 μ L of an antibody solution prepared at different concentrations was added, and reacted on ice for 15 minutes. Subsequently, 100 μ L of the mouse bone marrow derived effector cell solution (5×10^5 cells/well) was added, and cells were cultured in a 5% carbon dioxide gas incubator at 37°C for 4 hours. After the cultivation, the plate was centrifuged, and the radioactivity in 100 μ L of the culture supernatant was measured using a gamma counter. The specific chromium release rate was obtained by the following formula. Specific chromium release rate (%) = $(A-C) \times 100 / (B-C)$ "A" represents the mean value of the radioactivities (cpm) in each well, "B" represents the mean value of the radioactivities (cpm) in the wells in which 100 μ L of 2% NP-40 aqueous solution (Nonidet P-40, Code No. 252-23, Nacalai Tesque) and 50 μ L of 10% FBS/RPMI medium were added to the target cell, and "C" represents the mean value of the radioactivities (cpm) in the wells in which 150 μ L of 10% FBS/RPMI medium was added to the target cell. The test was carried out in triplicate and the mean value and the standard deviation were calculated for ADCC activity (%).

[0175] The results are shown in Fig. 9. It was revealed that GC33 antibody shows an ADCC activity when the concentration of antibody is 0.1 μ g/mL or higher, and shows stronger activity than GC199 antibody.

Example 20

Antitumor activity of GC33 antibody to mouse model transplanted with human hepatoma

20.1 Preparation of mouse model transplanted with human hepatoma

[0176] A human hepatoma cell line, HuH-7, was prepared at 5×10^7 cells/mL in a solution containing DMEM medium and MATRIGEL (BD Bioscience) at a ratio of 1:1. On the previous day, 100 μ L of an anti-asialo GM1 antibody solution (Wako Pure Chemicals, one vial was dissolved with 1 mL of distilled water for injection then added 4 mL of physiologic saline) was intraperitoneally administered to a SCID mouse (male, 5 weeks old, CLEA Japan, Inc.). The mouse was transplanted with 100 μ L of the above-mentioned cell suspension (5×10^6 cells/mouse) subcutaneously in the abdominal area.

20.2 Preparation and administration of antibody

[0177] Starting from the day 20 after the cell transplantation, an antibody solution prepared on the day of administration at 0.5 mg/mL (group of administration of 5 mg/kg) and at 0.1 mg/mL (group of administration of 1 mg/kg) with PBS(-) was administered to the mouse model transplanted with a human hepatoma cells at 10 mL/kg via the tail vein once a week for 3 weeks. As a negative control, PBS(-) (vehicle) was administered at 10 mL/kg via the tail vein once a week for 3 weeks in a similar manner. Both groups consisted of 6 mice each.

20.3 Evaluation of antitumor effect

[0178] The antitumor effect of GC33 antibody on the mouse model transplanted with human hepatoma cells was evaluated with the change in tumor volume with time and tumor weight at 1 week after the final administration. The tumor volume was calculated by the following formula.

$$\text{Tumor volume} = (\text{major axis}) \times (\text{minor axis}) \times (\text{minor axis}) / 2$$

[0179] As shown in Fig. 10, a significant inhibition of tumor growth was observed in the GC33 antibody group compared with the vehicle group.

[0180] Consequently, GC33 was shown to have an antitumor effect on the mouse model transplanted with a human hepatoma cells.

Example 21

Preparation of GC33 mouse-human chimeric antibody

[0181] The H chain and the L chain of GC33 were amplified by PCR using a synthetic oligonucleotide, which is complementary to the 5'-terminal nucleotide sequences and has a Kozak sequence and a HindIII site, and a synthetic oligonucleotide, which is complementary to the 3'-terminal nucleotide sequences and has a BamHI site. After digestion with HindIII and BamHI, the obtained PCR product was cloned into an expression vector, HEFgyl, in which a human IgG1 constant region was inserted, and an expression vector, HEFgk, in which a human kappa chain constant region was inserted (Sato et al., Mol Immunol. 1994; 371-381). The vectors were introduced into a CHO cell (DG44 cell line) in accordance with the above-mentioned method, and a stably expressing cell line was established. The antibody was purified from the culture supernatant using Hi Trap ProteinG HP (Amersham). The concentration of IgG in the culture supernatant was measured by a human IgG sandwich ELISA using goat anti-human IgG (BIOSOURCE) and goat anti-human IgG alkaline phosphatase conjugate (BIOSOURCE), and the concentration was determined by the comparison with a commercially available human IgG (Cappel).

Example 22

Measurement of CDC activity and ADCC activity using GC33 mouse-human chimeric antibody

[0182] In accordance with the methods described in Examples 16 and 17, the CDC activities and ADCC activities of GC33, M3C11 and M1E7 mouse-human chimeric antibodies were measured. As for the target cell, the CHO cell expressing full-length GPC3 was used for measuring the CDC activity and HepG2 was used for measuring the ADCC activity. The results are shown in Fig. 11 and Fig. 12, respectively. It was revealed that, in either test system, GC33 shows a strong CDC activity and ADCC activity compared with the other two antibodies.

Example 23

Epitope analysis for GC33

[0183] To determine the epitope for GC33 in detail, fusion proteins of a further shorter C-terminal peptide of GPC3 and GST were prepared, and analyzed by Western blotting. The prepared GPC3-derived peptide sequences contained in the GST-fusion protein are shown in Fig. 13. Since GC33 can bind to GC-4 (aa 537-563), but cannot bind to GC-5 (aa 550-563), it was considered that the epitope is located in a region containing at least part of the aa 537-550 region. First, the peptides GC-6 (G N S Q Q A T P K D N E I S (SEQ ID NO: 93)), GC-7 (G N S Q Q A T P (SEQ ID NO: 94)), GC-8 (Q Q A T P K D N (SEQ ID NO: 95)) and GC-9 (T P K D N E I S (SEQ ID NO: 96)) were prepared. A forward oligo DNA and a reverse oligo DNA were prepared which were designed in such a manner that the cleavage site of EcoRI recognition sequence is attached to the 5' end and the cleavage site of Sall recognition sequence is attached to the 3' end, respectively. The synthesis of the oligo DNAs was done by Espec Oligo Service. The DNA was purified with C-18 cartridge, phosphorylated at the 5' end and used for the analysis. Twenty-five microliters of the forward oligo DNA (10 μ M) and 25 μ L of the reverse oligo DNA (10 μ M) were mixed and reacted at 94 °C for 5 minutes, at 37 °C for 10 minutes, and at room temperature for 15 minutes, then left at 4 °C for 10 minutes to anneal the forward oligo DNA and the reverse oligo DNA. The concentration of the oligos was determined by the absorbance measurement at the molar ratio of the insert to the vector of 3:1. The oligos were cloned into EcoRI- and Sall-digested pGEX4T-3, and the nucleotide sequence was confirmed. A GST-fusion protein was prepared in accordance with the above-mentioned method, and purified using Glutathione Sepharose 4B. The purified proteins were separated by SDS-PAGE under reducing conditions, and analyzed by Western blotting using GC33. As a result, the antibody GC33 could not detect any GST-fusion protein strongly, suggesting that a longer sequence at the C-terminal side is needed for the binding of GC33 (Fig. 14). Based on the above prediction, GC-11 (A T P K D N E I S T (SEQ ID NO: 97)), GC-12 (P K D N E I S T F H (SEQ ID NO: 98)), GC-13 (D N E I S T F H N L (SEQ ID NO: 99)) and GC-14 (E I S T F H N L G N (SEQ ID NO: 100)) were prepared and evaluated in the same manner. As a result, GC-11, GC-12 and GC-13 bound to GC33 more strongly, suggesting that the epitope for GC33 is located in the sequence from 544th to 553rd (P K D N E I S T F H) at the C-terminus of GPC3.

Example 24

Humanization of GC33

[0184] Antibody sequence data were obtained from publicly disclosed Kabat Database (<ftp://ftp.ebi.ac.uk/pub/databases/kabat/>) and from ImMunoGeneTics Database (IMGT). The H chain variable region and the L chain variable region were separately subjected to homology search. As a result, the H chain variable region was found to have a high homology with DN13 (Smithson et al., Mol Immunol. 1999; 36: 113-124), and the L chain variable region was found to have a high homology with homo sapiens IGH mRNA for immunoglobulin kappa light chain VLJ region, partial cds, clone: K64 of the accession number of AB064105. The signal sequence of the accession number of S40357 that has a high homology with AB064105 was used as a signal sequence of the L chain. The complementarity determining region (hereinafter referred to as CDR) of GC33 were transplanted into the frame work regions (hereinafter referred to as FR) of these human antibodies to prepare a humanized antibody.

[0185] Specifically, synthetic oligo DNAs of approximately 50 bases were designed in such a manner that approximately 20 bases of them were hybridized and these synthetic oligo DNAs were assembled together by the PCR method to prepare genes encoding each of the variable regions. They were digested at the HindIII site inserted in the end of the 5'-terminal synthetic oligo DNA and the BamHI site inserted in the end of the 3' -terminal synthetic oligo DNA. The fragments were cloned into an expression vector, HEFgy1, in which a human IgG constant region was cloned, or an expression vector, HEFgx1, in which a human kappa chain constant region was cloned (Sato et. al., Mol Immunol. 1994; 371-381). The H chain and the L chain of the humanized GC33 constructed as above were named ver.a, respectively. The binding activity of the humanized GC33, whose H chain and the L chain were both ver.a (ver.a/ver.a), was lower than that of an antibody with mouse GC33 variable regions (mouse/mouse). Antibodies were constructed in which the mouse GC33 sequence and the ver.a sequence were chimerically combined (mouse/ver.a, ver.a/mouse) with regard to the H chain and the L chain, and their binding activities were evaluated. As a result, a decrease in binding activity was observed in ver.a/mouse, indicating that the decrease in binding activity due to amino acid replacement was attributed to the H chain (Fig. 15). Then, modified H chains, ver.c, ver.f, ver.h, ver.i, ver.j, ver.k were prepared. All these humanized GC33 showed a binding activity equivalent to that of a chimeric antibody having the mouse GC33 variable region (Fig. 15). The nucleotide sequences of the humanized GC33 H chain variable regions, ver.a, ver.c, ver.f, ver.h, ver.i, ver.j, ver.k were shown in SEQ ID NOs: 77, 78, 79, 80, 81, 82 and 83, respectively, and the amino acid sequences thereof were shown in SEQ ID NOs: 84, 85, 86, 87, 88, 89 and 90, respectively. The nucleotide sequence and the amino acid sequence of a humanized GC33 L chain variable region, ver.a are shown in SEQ ID NOs: 91 and 92, respectively. In humanized GC33 H chain variable regions ver.i, ver.j and ver.k, the 6th glutamic acid residue is replaced with glutamine residue. The heat stability of these antibodies was significantly increased.

Example 25

Modification of humanized GC33 L chain

[0186] As for the deamidation of protein, the reaction rate constant of deamidation is known to be dependent on the primary sequence. It is also known that Asn-Gly is particularly susceptible to deamidation (Rocinson et. al., Proc. Natl. Acad. Sci. USA 2001; 98: 944-949). As for Asn33 in the CDR1 of a humanized GC33 L chain ver.a shown in SEQ ID NO: 91, the primary sequence is Asn-Gly, which is predicted to be very susceptible to deamidation.

[0187] To evaluate the effect of deamidation of Asn33 on the binding activity, a modified antibody was prepared in which Asn33 was replaced with Asp. A point mutation was introduced using a Quick Change Site-Directed Mutagenesis Kit (Stratagene) was used. More specifically, 50 μ L of a reaction mixture containing 125 ng of a sense primer (CTT GTA CAC AGT GAC GGA AAC ACC TAT: SEQ ID NO: 172), 125 ng of an antisense primer (ATA GGT GTT TCC GTC ACT GTG TAC AAG: SEQ ID NO: 173), 5 μ L of 10x reaction buffer, 1 μ L of dNTP mix, 10 ng of HEFgk into which a humanized GC33 L chain ver.a had been cloned and 1 μ L of Pfu Turbo DNA Polymerase was subjected to PCR of 12 cycles consisting of 95°C for 30 seconds, 55°C for 1 minute and 68°C for 9 minutes. Subsequently, a restriction enzyme, DpnI, was added and digestion was carried out at 37°C for 2 hours, and the digested product was introduced into XL1Blue competent cell attached to the kit, whereby a transformant was obtained. The variable region was cleaved out from the clone in which each mutation was properly introduced, and cloned into HEFgk again. It was introduced into a COS7 cell using Eugene 6 (Roche) together with HEFgy1, in which a humanized GC33 H chain ver.k had been cloned. The antibody transiently expressed in the cell was recovered from the culture supernatant. The concentration of antibody was determined by a sandwich ELISA using the anti-human IgG antibody. The binding activity of the modified antibody was evaluated by an ELISA using an immunoplate coated with the soluble form of GPC3 core protein. As shown in Fig. 18, the binding activity was lost in the modified antibody (N33D) in which Asn33 had been replaced with Asp, suggesting that the effect of the deamidation of Asn33 on the binding activity was significant.

[0188] As a method of suppressing deamidation of Asn33, replacement of Gly34 with another amino acid has been reported (International Patent Application WO 03057881A1). In accordance with the above-mentioned method, G34 was replaced with any of 17 amino acids other than Cys and Met using a Quick Change Site-Directed Mutagenesis Kit to prepare a series of modified antibodies, namely, G34A, G34D, G34E, G34F, G34H, G34N, G34P, G34Q, G34I, G34K, G34L, G34V, G34W, G34Y, G34R, G34S and G34T. These antibodies were transiently expressed in COS7 cells, and the binding activity was evaluated using the culture supernatant. It was found that the binding activity is maintained even if G34 is replaced with another amino acid, except for Pro (G34P) and Val (G34V).

[0189] The amino acid sequences of the light chain CDR1 of the above-mentioned modified antibodies are shown in SEQ ID NO: 174 (G34A), SEQ ID NO: 175 (G34D), SEQ ID NO: 176 (G34E), SEQ ID NO: 177 (G34F), SEQ ID NO: 178 (G34H), SEQ ID NO: 179 (G34N), SEQ ID NO: 180 (G34T), SEQ ID NO: 181 (G34Q), SEQ ID NO: 182 (G34I), SEQ ID NO: 183 (G34K), SEQ ID NO: 184 (G34L), SEQ ID NO: 185 (G34S), SEQ ID NO: 186 (G34W), SEQ ID NO: 187 (G34Y), SEQ ID NO: 188 (G34R), SEQ ID NO: 189 (G34V) and SEQ ID NO: 190 (G34P), respectively. The amino acid sequences of the light chain variable regions of the above-mentioned modified antibodies are shown in SEQ ID NO: 191 (G34A), SEQ ID NO: 192 (G34D), SEQ ID NO: 193 (G34E), SEQ ID NO: 194 (G34F), SEQ ID NO: 195 (G34H), SEQ ID NO: 196 (G34N), SEQ ID NO: 197 (G34T), SEQ ID NO: 198 (G34Q), SEQ ID NO: 199 (G34I), SEQ ID NO: 200 (G34K), SEQ ID NO: 201 (G34L), SEQ ID NO: 202 (G34S), SEQ ID NO: 203 (G34W), SEQ ID NO: 204 (G34Y), SEQ ID NO: 205 (G34R), SEQ ID NO: 206 (G34V) and SEQ ID NO: 207 (G34P), respectively.

[0190] The antibody of the present invention can be used as a cell growth inhibitor, an anticancer agent or an agent for diagnosis of cancers.

Example 26

Preparation of human hepatoma cell line (SK-03) expressing full-length human GPC3

[0191] To obtain a cell line for evaluating a biological activity of the anti-GPC3 antibodies, a human hepatoma cell line expressing full-length GPC3 was established.

[0192] One microgram of a full-length human GPC3 gene expression vector treated with Pvu I was mixed with 2 μ L of FuGENE (Roche) to allow for complex formation. The complex was added to SK-HEP-1 cells (purchased from ATCC) for gene introduction. After incubation in CO₂ incubator for 24 hours, GPC3 expressing cells were selected using Dulbecco's MEM (D-MEM, SIGMA) containing Geneticin at a final concentration of 1 mg/mL and 10% FBS. The resulting Geneticin-resistant colonies were collected and cell cloning was carried out by the limiting dilution method. The expression of human GPC3 of each cell clone was assayed by flow cytometry using the chimeric antibody GC33 and FITC-labeled goat anti-human IgG antibody (ICN). In this way, a stably expressing cell line SK-03 was obtained.

Example 27

Comparison of CDC activity and ADCC activity of mouse-human chimeric antibodies

[0193] In order to directly compare the CDC activity and ADCC activity of the mouse-human chimeric antibodies GC33, M3C11, and M1E7 described in Example 22, the CDC activity and ADCC activity of three antibodies were measured in the same test system according to the method described in Examples 16 and 17. As for the target cell, the CHO cell expressing full-length GPC3 was used for measuring the CDC activity and SK-03 was used for measuring the ADCC activity. The results are shown in Fig. 19 and Fig. 20, respectively. It was revealed that, in either test system, GC33 shows a stronger CDC activity and ADCC activity compared with the other two antibodies.

Industrial Applicability

[0194] The antibody of the present invention can be used as a cell growth inhibitor, an anticancer agent and an agent for diagnosis of cancers.

SEQUENCE LISTING

<110> Chugai Seiyaku Kabushiki Kaisha
 <120> Anti-Glypican 3 Antibodies
 5 <130> N96765 GCW
 <150> JP 2004-203637
 <151> 2004-07-09
 10 <160> 173
 <170> PatentIn version 3.1
 <210> 1
 <211> 31
 15 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer
 20 <400> 1
 gatatcatgg ccgggaccgt gcgcaccgcg t 31
 <210> 2
 <211> 31
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer
 30 <400> 2
 gctagctcag tgcaccagga agaagaagca c 31
 <210> 3
 <211> 1743
 35 <212> DNA
 <213> homo sapiens
 <400> 3
 40 atggccggga ccgtgcgcac cgcgtgcttg gtggtggcga tgctgctcag cttggacttc 60
 ccgggacagg cgcagccccc gccgcgccg ccggacgcc cctgtcacca agtccgctcc 120
 ttcttccaga gactgcagcc cggactcaag tgggtgccag aaactcccgt gccaggatca 180
 gatttgcaag tatgtctccc taagggccca acatgctgct caagaaagat ggaagaaaaa 240
 45 taccaactaa cagcagcatt gaacatggaa cagctgcttc agtctgcaag tatggagctc 300
 aagttcttaa ttattcagaa tgctgcgggtt ttccaagagg cctttgaaat tgttgttcgc 360
 catgccaaga actacaccaa tgccatgttc aagaacaact acccaagcct gactccacaa 420
 gcttttgagt ttgtgggtga atttttcaca gatgtgtctc tctacatctt gggttctgac 480
 50 atcaatgtag atgacatggt caatgaattg tttgacagcc tgtttccagt catctataacc 540
 cagctaataa acccaggcct gcctgattca gccttgga tcaatgagt cctccgagga 600
 gcaagacgtg acctgaaagt atttgggaat ttccccaagc ttattatgac ccagggttcc 660
 aagtcactgc aagtcactag gatcttcctt caggctctga atcttggaat tgaagtgatc 720
 55 aacacaactg atcacctgaa gttcagtaag gactgtggcc gaatgctcac cagaatgtgg 780

tactgtctctt actgccaggg actgatgatg gttaaaccct gtggcgggta ctgcaatgtg 840
 gtcattgcaag gctgtatggc aggtgtggtg gagattgaca agtactggag agaatacatt 900
 5 ctgtcccttg aagaacttgt gaatggcatg tacagaatct atgacatgga gaacgtactg 960
 cttgggtctct tttcaacaat ccatgattct atccagtatg tccagaagaa tgcaggaaaag 1020
 ctgaccacca ctattggcaa gttatgtgcc cattctcaac aacgccaata tagatctgct 1080
 tattatcctg aagatctctt tattgacaag aaagtattaa aagttgctca ttagaacaat 1140
 10 gaagaaacct tatccagccg aagaagggaa ctaattcaga agttgaagtc tttcatcagc 1200
 ttctatagtg ctttgccctgg ctacatctgc agccatagcc ctgtggcgga aaacgacacc 1260
 ctttgctgga atggacaaga actcgtggag agatacagcc aaaaggcagc aaggaatgga 1320
 atgaaaaacc agttcaatct ccatgagctg aaaatgaagg gccctgagcc agtggtcagt 1380
 15 caaattattg acaaactgaa gcacattaac cagctcctga gaaccatgtc tatgccccaa 1440
 ggtagagtgc tggataaaaa cctggatgag gaagggtttg aaagtggaga ctgcggtgat 1500
 gatgaagatg agtgcattgg aggctctggt gatggaatga taaaagtga gaatcagctc 1560
 cgcttccttg cagaactggc ctatgatctg gatgtggatg atgcgcctgg aaacagtcag 1620
 20 caggcaactc cgaaggacaa cgagataagc acctttcaca acctcgggaa cgttcattcc 1680
 ccgtgaagc ttctcaccag catggccatc tcggtggtgt gcttcttctt cctggtgcac 1740
 tga
 <210> 4
 25 <211> 580
 <212> PRT
 <213> homo sapiens
 <400> 4
 30 Met Ala Gly Thr Val Arg Thr Ala Cys Leu Val Val Ala Met Leu Leu
 1 5 10 15
 Ser Leu Asp Phe Pro Gly Gln Ala Gln Pro Pro Pro Pro Pro Pro Asp
 20 25 30
 35 Ala Thr Cys His Gln Val Arg Ser Phe Phe Gln Arg Leu Gln Pro Gly
 35 40 45
 Leu Lys Trp Val Pro Glu Thr Pro Val Pro Gly Ser Asp Leu Gln Val
 50 55 60
 40 Cys Leu Pro Lys Gly Pro Thr Cys Cys Ser Arg Lys Met Glu Glu Lys
 65 70 75 80
 Tyr Gln Leu Thr Ala Arg Leu Asn Met Glu Gln Leu Leu Gln Ser Ala
 85 90 95
 45 Ser Met Glu Leu Lys Phe Leu Ile Ile Gln Asn Ala Ala Val Phe Gln
 100 105 110
 Glu Ala Phe Glu Ile Val Val Arg His Ala Lys Asn Tyr Thr Asn Ala
 115 120 125
 50 Met Phe Lys Asn Asn Tyr Pro Ser Leu Thr Pro Gln Ala Phe Glu Phe
 130 135 140
 Val Gly Glu Phe Phe Thr Asp Val Ser Leu Tyr Ile Leu Gly Ser Asp
 55 145 150 155 160

EP 1 674 111 A1

Ile Asn Val Asp Asp Met Val Asn Glu Leu Phe Asp Ser Leu Phe Pro
165 170 175
5 Val Ile Tyr Thr Gln Leu Met Asn Pro Gly Leu Pro Asp Ser Ala Leu
180 185 190
Asp Ile Asn Glu Cys Leu Arg Gly Ala Arg Arg Asp Leu Lys Val Phe
195 200 205
10 Gly Asn Phe Pro Lys Leu Ile Met Thr Gln Val Ser Lys Ser Leu Gln
210 215 220
Val Thr Arg Ile Phe Leu Gln Ala Leu Asn Leu Gly Ile Glu Val Ile
225 230 235 240
15 Asn Thr Thr Asp His Leu Lys Phe Ser Lys Asp Cys Gly Arg Met Leu
245 250 255
Thr Arg Met Trp Tyr Cys Ser Tyr Cys Gln Gly Leu Met Met Val Lys
260 265 270
20 Pro Cys Gly Gly Tyr Cys Asn Val Val Met Gln Gly Cys Met Ala Gly
275 280 285
Val Val Glu Ile Asp Lys Tyr Trp Arg Glu Tyr Ile Leu Ser Leu Glu
290 295 300
25 Glu Leu Val Asn Gly Met Tyr Arg Ile Tyr Asp Met Glu Asn Val Leu
305 310 315 320
Leu Gly Leu Phe Ser Thr Ile His Asp Ser Ile Gln Tyr Val Gln Lys
325 330 335
30 Asn Ala Gly Lys Leu Thr Thr Thr Ile Gly Lys Leu Cys Ala His Ser
340 345 350
Gln Gln Arg Gln Tyr Arg Ser Ala Tyr Tyr Pro Glu Asp Leu Phe Ile
355 360 365
35 Asp Lys Lys Val Leu Lys Val Ala His Val Glu His Glu Glu Thr Leu
370 375 380
Ser Ser Arg Arg Arg Glu Leu Ile Gln Lys Leu Lys Ser Phe Ile Ser
385 390 395 400
40 Phe Tyr Ser Ala Leu Pro Gly Tyr Ile Cys Ser His Ser Pro Val Ala
405 410 415
Glu Asn Asp Thr Leu Cys Trp Asn Gly Gln Glu Leu Val Glu Arg Tyr
420 425 430
45 Ser Gln Lys Ala Ala Arg Asn Gly Met Lys Asn Gln Phe Asn Leu His
435 440 445
Glu Leu Lys Met Lys Gly Pro Glu Pro Val Val Ser Gln Ile Ile Asp
450 455 460
50 Lys Leu Lys His Ile Asn Gln Leu Leu Arg Thr Met Ser Met Pro Lys
465 470 475 480
Gly Arg Val Leu Asp Lys Asn Leu Asp Glu Glu Gly Phe Glu Ser Gly
485 490 495
55

EP 1 674 111 A1

Asp Cys Gly Asp Asp Glu Asp Glu Cys Ile Gly Gly Ser Gly Asp Gly
500 505 510
5 Met Ile Lys Val Lys Asn Gln Leu Arg Phe Leu Ala Glu Leu Ala Tyr
515 520 525
Asp Leu Asp Val Asp Asp Ala Pro Gly Asn Ser Gln Gln Ala Thr Pro
530 535 540
10 Lys Asp Asn Glu Ile Ser Thr Phe His Asn Leu Gly Asn Val His Ser
545 550 555 560
Pro Leu Lys Leu Leu Thr Ser Met Ala Ile Ser Val Val Cys Phe Phe
565 570 575
15 Phe Leu Val His
580
<210> 5
<211> 31
20 <212> DNA
<213> Artificial Sequence
<220>
<223> PCR primer
25 <400> 5
atagaattcc accatggccg ggaccgtgcg c 31
<210> 6
<211> 31
30 <212> DNA
<213> Artificial Sequence
<220>
<223> PCR primer
35 <400> 6
ataggatccc ttcagcgggg aatgaacgtt c 31
<210> 7
<211> 21
40 <212> DNA
<213> Artificial Sequence
<220>
<223> PCR primer
45 <400> 7
gggccagtgg atagacagat g 21
<210> 8
50 <211> 24
<212> DNA
<213> Artificial Sequence
<220>
55 <223> PCR primer

<400> 8
 cagggggccag tggatagacc gatg 24
 <210> 9
 5 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 10 <223> PCR primer
 <400> 9
 cagggggccag tggatagact gatg 24
 15 <210> 10
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 20 <220>
 <223> PCR primer
 <400> 10
 gctcactgga tgggtgggaag atg 23
 25 <210> 11
 <211> 1392
 <212> DNA
 <213> Mus musculus
 30 <400> 11
 atgaacttcg ggctcacctt gattttcctt gtccttactt taaaaggtgt ccagtgtgag 60
 gtgcaactgg tggagtctgg gggaggctta gtgaagcctg gaggatccct gaaactctcc 120
 tgtgcagcct ctggattcac tttcagtgc tatgccatgt cttgggttcg ccagattcca 180
 35 gagaagatac tggagtgggt cgcagccatt gatagtagtg gtggtgacac ctactattta 240
 gacactgtga aggaccgatt caccatctcc agagacaatg ccaataatac cctgcacctg 300
 caaatgcgca gtctgaggte tgaggacaca gccttgtatt actgtgtaag acaggggggg 360
 gcttactggg gccaaaggac tctgggtcact gtctctgcag ctagcaccaa gggcccatcg 420
 40 gtcttccccc tggcacctc ctccaagagc acctctgggg gcacagcggc cctgggctgc 480
 ctggtcaagg actacttccc cgaaccggtg acggtgtcgt ggaactcagg cgccctgacc 540
 agcggcgtgc acaccttccc ggctgtccta cagtcctcag gactctactc cctcagcagc 600
 45 gtggtgaccg tgccctccag cagcttgggc acccagacct acatctgcaa cgtgaatcac 660
 aagcccagca acaccaaggt ggacaagaaa gttgagccca aatcttgtga caaaactcac 720
 acatgccac cgtgccagc acctgaactc ctggggggac cgtcagtctt cctcttcccc 780
 ccaaaaccca aggacacct catgatctcc cggaccctg aggtcacatg cgtggtggtg 840
 50 gacgtgagcc acgaagacct tgaggtaag ttcaactggt acgtggacgg cgtggagggtg 900
 cataatgcca agacaaagcc gcgggaggag cagtacaaca gcacgtaccg tgtggtcagc 960
 gtcctcaccg tctgcacca ggactggctg aatggcaagg agtacaagtg caaggctctc 1020
 aacaaagccc tcccagcccc catcgagaaa accatctcca aagccaaagg gcagccccga 1080
 55 gaaccacagg tgtacacct gccccatcc cgggatgagc tgaccaagaa ccaggtcagc 1140

EP 1 674 111 A1

ctgacctgcc tgggtcaaagg cttctatccc agcgacatcg ccgtggagtg ggagagcaat 1200
 gggcagccgg agaacaacta caagaccacg cctcccgtgc tggactccga cggctccttc 1260
 5 ttcctctaca gcaagctcac cgtggacaag agcaggtggc agcaggggaa cgtcttctca 1320
 tgctccgtga tgcattgagc tctgcacaac cactacacgc agaagagcct ctccctgtct 1380
 ccgggtaaat ga 1392
 <210> 12
 10 <211> 342
 <212> DNA
 <213> Mus musculus
 <400> 12
 15 gaggtgcacc tgggtggagtc tgggggaggc ttagtgaagc ctggagggtc cctgaaactc 60
 tcctgtgcag cctctggatt cactttcagt aactatgcca tgtcttgggt tcgccagact 120
 ccagagaaga ggctggagtg ggtcgcagcc attaataata atggtgatga cacctactat 180
 ttagacactg tgaaggaccg attcaccatc tccagagaca atgccaagaa caccctgtac 240
 20 ctgcaaatga gcagtctgag gtctgaggac acagccctgt attactgtgt aagacaaggg 300
 ggggcttact ggggcccaagg gactctggtc actgtctctg ca 342
 <210> 13
 <211> 1413
 25 <212> DNA
 <213> Mus musculus
 <400> 13
 atgggatgga actggatctt tattttaatc ctgtcagtaa ctacaggtgt ccaactotgag 60
 30 gtccagctgc agcagttctg acctgagctg gtgaagcctg gggcttcagt gaagatatcc 120
 tgcaaggctt ctggttactc attcactggc tactacatgc actgggtgaa gcaaagtcct 180
 gaaaagagcc ttgagtggat tggagagatt aatcctagca ctggtggtag tacctacaac 240
 35 cagaagttca aggccaaggc cacattgact gtagacaaat cctccagcac agcctacatg 300
 cagctcaaga gcttgacatc tgaggactct gcagtctatt actgtgcaag gaggggcgga 360
 ttaactggga cgagcttctt tgttactgg ggccaaggga ctctggtcac tgtctctgca 420
 gctagcacca agggcccatc ggtcttcccc ctggcaccct cctccaagag cactcttggg 480
 40 ggcacagcgg ccttgggctg cctggtcaag gactacttcc ccgaaccggg gacggtgtcg 540
 tggaactcag gcgccctgac cagcggcgtg cacaccttcc cggctgtcct acagtcctca 600
 ggactctact cctcagcag cgtggtgacc gtgccctcca gcagcttggg caccagacc 660
 tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc 720
 45 aaatcttgtg acaaaactca cacatgcca cgtgcccag cacctgaact cctgggggga 780
 ccgtcagttt tctcttccc cccaaaaccc aaggacacc tcattgatct ccggaccctt 840
 gaggtcacat gcgtggtggt ggacgtgagc caggaagacc ctgaggtcaa gttcaactgg 900
 tacgtggacg gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac 960
 50 agcacgtacc gtgtggtcag cgtcctcacc gtctgcacc aggactggct gaatggcaag 1020
 gagtacaagt gcaaggtctc caacaaagcc ctcccagccc ccatogagaa aaccatctcc 1080
 aaagccaaag ggcagccccg agaaccacag gtgtacacccc tgcccccatc ccgggatgag 1140
 ctgaccaaga accaggtcag cctgacctgc ctgggtcaaag gcttctatcc cagcgacatc 1200
 55 gccgtggagt gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccgtg 1260

EP 1 674 111 A1

ctggactcgc acggetcctt ctctctctac agcaagctca ccgtggacaa gagcaggtgg 1320
 cagcagggga acgtcttctc atgctccgtg atgcatgagg ctctgcacaa ccactacacg 1380
 5 cagaagagcc tctccctgtc tccgggtaaa tga 1413
 <210> 14
 <211> 354
 <212> DNA
 10 <213> Mus musculus
 <400> 14
 caggtcactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
 acttgttctt tctctgggtt ttactgagc atttatggta tgggtgtagg ttggattcgt 120
 15 cagccttcag ggatgggtct ggagtggctg gccaacattt ggtggtatga tgctaagtac 180
 tataactctg acctgaagag ccggctcaca atctccaagg atacctcaa caaccaggtg 240
 ttctcaaga tctccagtgt ggacacttca gatactgcc catactactg tgctcaaagt 300
 ggactggcct gggttgctta ctggggccaa gggactctgg tcaactgtctc tgca 354
 20 <210> 15
 <211> 354
 <212> DNA
 <213> Mus musculus
 25 <400> 15
 caggtcactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
 acttgttctt tctctgggtt ttactgagc atttatggta tgggtgtagg ttggattcgt 120
 cagccttcag ggaagggctt ggagtggctg gccaacattt ggtggaatga tgataagtac 180
 30 tataactcag ccctgaagag ccggctcaca atctccaagg atacctcaa caaccaggta 240
 ttctcaaga tctccagtgt ggacactgca gatactgcc catactactg tgctcaaata 300
 gggtacttct actttgacta ctggggccaa ggcaccactc tcacagtctc ctca 354
 35 <210> 16
 <211> 1416
 <212> DNA
 <213> Mus musculus
 40 <400> 16
 atgaacttcg ggctcacctt gattttctct gtcttactt taaaagggtg ccagtgtgag 60
 gtgcagctgg tggagtctgg gggagactta gtgaagcctg gagggaccct gaaactctcc 120
 tgtgcagcct ctggatccac ttctagtaac tatgccatgt cttgggttcg ccagactcca 180
 45 gagaagaggc tggagtgggt cgcagccatt gatagtaat gaggtaccac ctactatcca 240
 gacactatga aggaccgatt caccatttcc agagacaatg ccaagaacac cctgtacctg 300
 caaatgaaca gtctgaggtc tgaagacaca gccttttatc actgtacaag acataatgga 360
 gggatatgaaa actacggctg gtttgcttac tggggccaag ggactctggt cactgtctct 420
 50 gcagctagca ccaagggccc atcgggtctt cccctggcac cctcctcaa gagcacctct 480
 gggggcacag cggccctggg ctgcctggtc aaggactact tccccgaacc ggtgacggtg 540
 tcgtggaact caggcgccct gaccagcggc gtgcacacct tcccggctgt cctacagtcc 600
 tcaggactct actccctcag cagcgtggtg accgtgccct ccagcagctt gggcaccacg 660
 55 acctacatct gcaacgtgaa tcacaagccc agcaacacca aggtggacaa gaaagttgag 720

EP 1 674 111 A1

	cccaaattctt	gtgacaaaac	tcacacatgc	ccaccgtgcc	cagcacctga	actcctgggg	780
	ggaccgtcag	tcttcctctt	cccccaaaa	cccaaggaca	ccctcatgat	ctcccgacc	840
5	cctgaggtca	catgcgtggt	ggtggacgtg	agccacgaag	accctgaggt	caagttcaac	900
	tggtacgtgg	acggcggtga	ggtgcataat	gccaaagaaa	agccgcggga	ggagcagtag	960
	aacagcacgt	accgtgtggt	cagcgtcctc	accgtcctgc	accaggactg	gctgaatggc	1020
	aaggagtaca	agtgcaaggt	ctccaacaaa	gccctcccag	cccccatcga	gaaaaccatc	1080
10	tccaaagcca	aagggcagcc	ccgagaacca	caggtgtaca	ccctgcccc	atcccgggat	1140
	gagctgacca	agaaccaggt	cagcctgacc	tgctgtgtca	aaggcttcta	tcccagcgac	1200
	atcgccgtgg	agtgggagag	caatgggcag	ccggagaaca	actacaagac	cacgcctccc	1260
	gtgctggact	ccgacggctc	cttcttcctc	tacagcaagc	tcaccgtgga	caagagcagg	1320
15	tggcagcagg	ggaacgtctt	ctcatgctcc	gtgatgcatg	aggctctgca	caaccactac	1380
	acgcagaaga	gcctctccct	gtctccgggt	aaatga			1416
	<210>	17					
	<211>	366					
20	<212>	DNA					
	<213>	Mus musculus					
	<400>	17					
	gaggtgcagc	tggtggagtc	tgggggagac	ttagtgaagc	ctggagggtc	cctgaaactc	60
25	tcctgtgcag	cctctggatt	cactttcagt	agctatgcca	tgtcttgggt	tcgccagact	120
	ccagagaaga	ggctggagtg	ggtcgcagcc	attaatagta	atggaggtag	cacctactat	180
	ccagacacta	tgaaggaccg	attcaccatc	tccagagaca	atgccaaaga	cacctgtac	240
	ctgcaaatga	gcagtctgag	gtctgaagac	tcagccttgt	attactgtac	aagacataat	300
30	ggagggtatg	aaaactacgg	ctgggttgct	tactggggcc	aagggactct	ggctcaactgc	360
	tctgca					366	
	<210>	18					
	<211>	1413					
35	<212>	DNA					
	<213>	Mus musculus					
	<400>	18					
40	atggaatcta	actggatact	tccttttatt	ctgtcggtag	cttcaggggt	ctactcagag	60
	gttcagctcc	agcagtctgg	gactgtgctg	gcaaggcctg	gggcttcagt	gaagatgtcc	120
	tgcaaggctt	ctggctacac	ctttactggc	tactggatgc	gctgggtaaa	acagaggcct	180
	ggacagggtc	tggaatggat	tggcgctatt	tatcctggaa	atagtgtata	aacatacaac	240
45	cagaagttca	agggcaaggc	caaactgact	gcagtcacat	ctgtcagcac	tgctacatg	300
	gaactcagca	gcctgacaaa	tgaggactct	gcggtctatt	actgttcaag	atcgggggac	360
	ctaactgggg	ggtttgctta	ctggggccaa	gggactctgg	tcactgtctc	tacagccaaa	420
	gctagcacca	agggcccatc	ggtcttcccc	ctggcaccct	cctccaagag	cacctctggg	480
50	ggcacagcgg	ccctgggctg	cctgggtcaag	gactacttcc	ccgaaccggg	gacggtgtcg	540
	tggaactcag	gcgccctgac	cagcggcgtg	cacaccttcc	cggtgtcct	acagtcctca	600
	ggactctact	ccctcagcag	cgtgggtgacc	gtgccctcca	gcagcttggg	caccagacc	660
	tacatctgca	acgtgaatca	caagcccagc	aacaccaagg	tggacaagaa	agttgagccc	720
55	aaatcttgtg	acaaaactca	cacatgccca	ccgtgcccag	cacctgaact	cctgggggga	780

ccgtcagtct tctcttcccc cccaaaaccc aaggacaccc tcatgatctc ccggaccacct 840
 gaggtcacat gcgtgggtgt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg 900
 5 tacgtggacg gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac 960
 agcacgtacc gtgtggtcag cgtcctcacc gtccctgcacc aggactggct gaatggcaag 1020
 gagtacaagt gcaaggtctc caacaaagcc ctcccagccc ccatcgagaa aaccatctcc 1080
 aaagccaaag ggcagccccg agaaccacag gtgtacaccc tgcccccatc ccgggatgag 1140
 10 ctgaccaaga accagggtcag cctgacctgc ctgggtcaaag gcttctatcc cagcgacatc 1200
 gccgtggagt gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccgtg 1260
 ctggactccg acggctcctt ctctccttac agcaagctca ccgtggacaa gagcaggtgg 1320
 cagcagggga acgtcttctc atgctccgtg atgcatgagg ctctgcacaa ccactacacg 1380
 15 cagaagagcc tctccctgtc tccgggtaaa tga 1413
 <210> 19
 <211> 357
 <212> DNA
 20 <213> Mus musculus
 <400> 19
 gaggttcagc tccagcagtc tgggactgtg ctggcaaggc ctggggcttc agtgaagatg 60
 tcttgcaagg cttctggcta cacctttacc ggctactgga tgcactgggt aaaacagagg 120
 25 cctggacagg gtctggaatg gattggcgct atttatcctg gaaatagtga tactaactac 180
 aaccagaagt tcaagggcaa ggccaaactg actgcagtca catctgccag cactgcctac 240
 atggagctca gcagcctgac aaatgaggac gctgcggtct atcactgtac aagatcgggg 300
 gacctaactg gggggcttgc ttactggggc caagggactc tggtcactgt ctctgca 357
 30 <210> 20
 <211> 372
 <212> DNA
 35 <213> Mus musculus
 <400> 20
 cagggtccagc tgcagcagcc tggggctgaa ctggtgaagc ctggggcttc agtgaaactg 60
 tcttgcaagg cttctggata caccttact agctactgga tgcattgggt gaagcagagg 120
 40 cctggacaag gccttgagtg gatcgagag attgatcctt ctgatagtta tacttactac 180
 aatcaaaagt tcaggggcaa ggccacattg actgtagaca aatcctccaa cacagcctac 240
 atgcaactca gcagcctgac atctgaggac tctgcggtct attactgttc aagatcaaact 300
 ctgggtgatg gtcactaccg gtttctgcg tttccttact ggggccaaag gactctggtc 360
 45 actgtctctg ca 372
 <210> 21
 <211> 372
 <212> DNA
 50 <213> Mus musculus
 <400> 21
 cagggtccaac tgcagcagcc tggggctgaa ctggtgaaac ctggggcttc agtgaagctg 60
 tcttgcaagg cttctggcta caccttcacc agctactgga tgcactgggt gaaacagagg 120
 55 cctggacaag gccttgaatg gattggtaca attgaccctt ctgatagtga aactcactac 180

aatctacagt tcaaggacac ggccacattg actgtagaca aatcctccag cacagcctac 240
 atgcagctca gcagcctgac atctgaggac tctgcggtct attattgtat aagaggcgcc 300
 5 ttctatagtt cctatagtta ctgggcctgg tttgcttact ggggccaagg gactctggtc 360
 actgtctctg ca 372
 <210> 22
 <211> 463
 10 <212> PRT
 <213> Mus musculus
 <400> 22
 Met Asn Phe Gly Leu Thr Leu Ile Phe Leu Val Leu Thr Leu Lys Gly
 15 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys
 20 20 25 30
 Pro Gly Gly Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
 25 35 40 45
 Ser Arg Tyr Ala Met Ser Trp Val Arg Gln Ile Pro Glu Lys Ile Leu
 50 55 60
 Glu Trp Val Ala Ala Ile Asp Ser Ser Gly Gly Asp Thr Tyr Tyr Leu
 25 65 70 75 80
 Asp Thr Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asn Ala Asn Asn
 85 90 95
 Thr Leu His Leu Gln Met Arg Ser Leu Arg Ser Glu Asp Thr Ala Leu
 30 100 105 110
 Tyr Tyr Cys Val Arg Gln Gly Gly Ala Tyr Trp Gly Gln Gly Thr Leu
 115 120 125
 Val Thr Val Ser Ala Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 35 130 135 140
 Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 145 150 155 160
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 40 165 170 175
 Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 180 185 190
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 45 195 200 205
 Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
 210 215 220
 Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His
 50 225 230 235 240
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 245 250 255
 55 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr

EP 1 674 111 A1

260 265 270
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 275 280 285
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 290 295 300
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 305 310 315 320
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 325 330 335
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 340 345 350
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 355 360 365
 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 370 375 380
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 385 390 395 400
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 405 410 415
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 420 425 430
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 435 440 445
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 450 455 460
 <210> 23
 <211> 114
 <212> PRT
 <213> Mus musculus
 <400> 23
 Glu Val His Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
 35 40 45
 Ala Ala Ile Asn Asn Asn Gly Asp Asp Thr Tyr Tyr Leu Asp Thr Val
 50 55 60
 Lys Asp Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Ser Ser Leu Arg Ser Glu Asp Thr Ala Leu Tyr Tyr Cys
 85 90 95

EP 1 674 111 A1

Val Arg Gln Gly Gly Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
100 105 110

Ser Ala

<210> 24

<211> 470

<212> PRT

<213> Mus musculus

<400> 24

Met Gly Trp Asn Trp Ile Phe Ile Leu Ile Leu Ser Val Thr Thr Gly
1 5 10 15

Val His Ser Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys
20 25 30

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe
35 40 45

Thr Gly Tyr Tyr Met His Trp Val Lys Gln Ser Pro Glu Lys Ser Leu
50 55 60

Glu Trp Ile Gly Glu Ile Asn Pro Ser Thr Gly Gly Thr Thr Tyr Asn
65 70 75 80

Gln Lys Phe Lys Ala Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser
85 90 95

Thr Ala Tyr Met Gln Leu Lys Ser Leu Thr Ser Glu Asp Ser Ala Val
100 105 110

Tyr Tyr Cys Ala Arg Arg Gly Gly Leu Thr Gly Thr Ser Phe Phe Ala
115 120 125

Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Ala Ser Thr Lys
130 135 140

Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly
145 150 155 160

Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
165 170 175

Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
180 185 190

Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
195 200 205

Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn
210 215 220

Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro
225 230 235 240

Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
245 250 255

Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp

EP 1 674 111 A1

260 265 270
 Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
 275 280 285
 5 Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
 290 295 300
 Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
 10 305 310 315 320
 Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
 325 330 335
 Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro
 15 340 345 350
 Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
 355 360 365
 Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn
 20 370 375 380
 Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
 385 390 395 400
 Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
 25 405 410 415
 Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
 420 425 430
 Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
 30 435 440 445
 Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
 450 455 460
 Ser Leu Ser Pro Gly Lys
 35 465 470
 <210> 25
 <211> 118
 40 <212> PRT
 <213> Mus musculus
 <400> 25
 Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
 45 1 5 10 15
 Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Tyr
 20 25 30
 Gly Met Gly Val Gly Trp Ile Arg Gln Pro Ser Gly Met Gly Leu Glu
 50 35 40 45
 Trp Leu Ala Asn Ile Trp Trp Tyr Asp Ala Lys Tyr Tyr Asn Ser Asp
 50 55 60
 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Asn Asn Gln Val
 55 65 70 75 80

EP 1 674 111 A1

Phe Leu Lys Ile Ser Ser Val Asp Thr Ser Asp Thr Ala Thr Tyr Tyr
 85 90 95
 5 Cys Ala Gln Met Gly Leu Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr
 100 105 110
 Leu Val Thr Val Ser Ala
 115
 10 <210> 26
 <211> 118
 <212> PRT
 <213> Mus musculus
 15 <400> 26
 Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
 1 5 10 15
 Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Ile Tyr
 20 20 25 30
 Gly Met Gly Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu
 35 40 45
 Trp Leu Ala Asn Ile Trp Trp Asn Asp Asp Lys Tyr Tyr Asn Ser Ala
 25 50 55 60
 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Asn Asn Gln Val
 65 70 75 80
 Phe Leu Lys Ile Ser Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr
 30 85 90 95
 Cys Ala Gln Ile Gly Tyr Phe Tyr Phe Asp Tyr Trp Gly Gln Gly Thr
 100 105 110
 Thr Leu Thr Val Ser Ser
 35 115
 <210> 27
 <211> 471
 <212> PRT
 40 <213> Mus musculus
 <400> 27
 Met Asn Phe Gly Leu Thr Leu Ile Phe Leu Val Leu Thr Leu Lys Gly
 45 1 5 10 15
 Val Gln Cys Glu Val Gln Leu Val Glu Ser Gly Gly Asp Leu Val Lys
 20 25 30
 Pro Gly Gly Thr Leu Lys Leu Ser Cys Ala Ala Ser Gly Ser Thr Phe
 35 40 45
 50 Ser Asn Tyr Ala Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu
 50 55 60
 Glu Trp Val Ala Ala Ile Asp Ser Asn Gly Gly Thr Thr Tyr Tyr Pro
 55 65 70 75 80

EP 1 674 111 A1

Asp Thr Met Lys Asp Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
 85 90 95
 5 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ser Glu Asp Thr Ala Phe
 100 105 110
 Tyr His Cys Thr Arg His Asn Gly Gly Tyr Glu Asn Tyr Gly Trp Phe
 115 120 125
 10 Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Ala Ser Thr
 130 135 140
 Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser
 145 150 155 160
 15 Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu
 165 170 175
 Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His
 180 185 190
 20 Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser
 195 200 205
 Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys
 210 215 220
 25 Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu
 225 230 235 240
 Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro
 245 250 255
 30 Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 260 265 270
 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 275 280 285
 35 Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
 290 295 300
 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
 305 310 315 320
 40 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 325 330 335
 Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu
 340 345 350
 45 Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 355 360 365
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys
 370 375 380
 50 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 385 390 395 400
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 405 410 415
 55

Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 420 425 430
 5 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
 435 440 445
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 450 455 460
 10 Leu Ser Leu Ser Pro Gly Lys
 465 470
 <210> 28
 <211> 122
 15 <212> PRT
 <213> Mus musculus
 <400> 28
 20 Glu Val Gln Leu Val Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Gly
 1 5 10 15
 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 25 Ala Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
 35 40 45
 Ala Ala Ile Asn Ser Asn Gly Gly Thr Thr Tyr Tyr Pro Asp Thr Met
 50 55 60
 30 Lys Asp Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Ser Ser Leu Arg Ser Glu Asp Ser Ala Leu Tyr Tyr Cys
 85 90 95
 35 Thr Arg His Asn Gly Gly Tyr Glu Asn Tyr Gly Trp Phe Ala Tyr Trp
 100 105 110
 Gly Gln Gly Thr Leu Val Thr Val Ser Ala
 115 120
 40 <210> 29
 <211> 470
 <212> PRT
 <213> Mus musculus
 45 <400> 29
 Met Glu Ser Asn Trp Ile Leu Pro Phe Ile Leu Ser Val Ala Ser Gly
 1 5 10 15
 50 Val Tyr Ser Glu Val Gln Leu Gln Gln Ser Gly Thr Val Leu Ala Arg
 20 25 30
 Pro Gly Ala Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35 40 45
 55 Thr Gly Tyr Trp Met Arg Trp Val Lys Gln Arg Pro Gly Gln Gly Leu
 50 55 60

EP 1 674 111 A1

	Glu Trp Ile Gly Ala Ile Tyr Pro Gly Asn Ser Asp Thr Thr Tyr Asn	
	65	70 75 80
5	Gln Lys Phe Lys Gly Lys Ala Lys Leu Thr Ala Val Thr Ser Val Ser	
		85 90 95
	Thr Ala Tyr Met Glu Leu Ser Ser Leu Thr Asn Glu Asp Ser Ala Val	
	100	105 110
10	Tyr Tyr Cys Ser Arg Ser Gly Asp Leu Thr Gly Gly Phe Ala Tyr Trp	
	115	120 125
	Gly Gln Gly Thr Leu Val Thr Val Ser Thr Ala Lys Ala Ser Thr Lys	
	130	135 140
15	Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly	
	145	150 155 160
	Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro	
	165	170 175
20	Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr	
	180	185 190
	Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val	
	195	200 205
25	Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn	
	210	215 220
	Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro	
	225	230 235 240
30	Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu	
	245	250 255
	Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp	
	260	265 270
35	Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp	
	275	280 285
	Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly	
	290	295 300
40	Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn	
	305	310 315 320
	Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp	
	325	330 335
45	Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro	
	340	345 350
	Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu	
	355	360 365
50	Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn	
	370	375 380
	Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile	
55	385	390 395 400

EP 1 674 111 A1

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
405 410 415
5 Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
420 425 430
Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
435 440 445
10 Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
450 455 460
Ser Leu Ser Pro Gly Lys
465 470
15 <210> 30
<211> 119
<212> PRT
<213> Mus musculus
20 <400> 30
Glu Val Gln Leu Gln Gln Ser Gly Thr Val Leu Ala Arg Pro Gly Ala
1 5 10 15
25 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30
Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45
30 Gly Ala Ile Tyr Pro Gly Asn Ser Asp Thr Asn Tyr Asn Gln Lys Phe
50 55 60
Lys Gly Lys Ala Lys Leu Thr Ala Val Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80
35 Met Glu Leu Ser Ser Leu Thr Asn Glu Asp Ala Ala Val Tyr His Cys
85 90 95
Thr Arg Ser Gly Asp Leu Thr Gly Gly Leu Ala Tyr Trp Gly Gln Gly
100 105 110
40 Thr Leu Val Thr Val Ser Ala
115
<210> 31
<211> 124
45 <212> PRT
<213> Mus musculus
<400> 31
Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
50 1 5 10 15
Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30
55 Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

EP 1 674 111 A1

Gly Glu Ile Asp Pro Ser Asp Ser Tyr Thr Tyr Tyr Asn Gln Lys Phe
 50 55 60
 5 Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Asn Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 10 Ser Arg Ser Asn Leu Gly Asp Gly His Tyr Arg Phe Pro Ala Phe Pro
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
 115 120
 15 <210> 32
 <211> 124
 <212> PRT
 <213> Mus musculus
 20 <400> 32
 Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 25 20 25 30
 Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Thr Ile Asp Pro Ser Asp Ser Glu Thr His Tyr Asn Leu Gln Phe
 30 50 55 60
 Lys Asp Thr Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 35 85 90 95
 Ile Arg Gly Ala Phe Tyr Ser Ser Tyr Ser Tyr Trp Ala Trp Phe Ala
 100 105 110
 Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
 40 115 120
 <210> 33
 <211> 717
 <212> DNA
 45 <213> Mus musculus
 <400> 33
 atgagtcctg cccagttcct gtttctgtta gtgctctgga ttcgggaaac caacggtgat 60
 50 gttgtgatga cccagactcc actcactttg tcggttacca ttggacaacc agcctccatc 120
 tcttgcaagt caagtcagag cctcttagat agtgatggaa agacatatat gaattgggtg 180
 ttacagaggc caggccagtc tccaaagcgc ctaatctatc tggtgtctaa attggactct 240
 ggagccccctg acagggttcac tggcagtgga tcaggacag atttcacact gaaaatcagt 300
 55 agagtggagg ctgaggattt gggaatttat tattgctggc aaggtacaca ttttccgctc 360

EP 1 674 111 A1

acgttcggtg ctgggaccaa gctggagctg aaacgtacgg tggctgcacc atctgtcttc 420
atcttcccg ccatctgatga gcagttgaaa tctggaactg cctctgttgt gtgcctgctg 480
5 aataacttct atcccagaga ggccaaagta cagtggaagg tggataacgc cctccaatcg 540
ggtaactccc aggagagtgt cacagagcag gacagcaagg acagcaccta cagcctcagc 600
agcaccctga cgctgagcaa agcagactac gagaaacaca aagtctacgc ctgcgaagtc 660
acccatcagg gcctgagctc gcccgtcaca aagagcttca acaggggaga gtgttga 717
10 <210> 34
<211> 336
<212> DNA
<213> Mus musculus
15 <400> 34
gatgttgtga tgaccagtc tccactcact ttgtcgatta ccattggaca accagcctcc 60
atctcttgca agtcaagtca gagcctotta gatagtgatg gaaagacata tttgaattgg 120
ttgttacaga ggccaggcca gtctocaaag cgcctaattct atctgggtgc taaactggac 180
20 tctggagtcc ctgacaggtt cactggcagt ggatcaggga cagatttctc actgaaaatc 240
agcagagtgg aggtgagga tttgggaatt tattattgct ggcaaggtag acattttcog 300
ctcacgttcg gtgctgggac caagctggag ctgaaa 336
<210> 35
25 <211> 717
<212> DNA
<213> Mus musculus
<400> 35
30 atgagtcctg tccagttcct gtttctgtta atgctctgga ttcaggaaac caacgggtgat 60
gttgtgatga cccagactcc actgtctttg tcggttacca ttggacaacc agcctctatc 120
tcttgcaagt caagtcagag cctcttatat agtaatggaa agacatattt gaattggtta 180
35 caacagaggc ctggccaggc tccaaagcac ctaatgtatc aggtgtccaa actggaccct 240
ggcatccctg acagggttcag tggcagtgga tcagaaacag attttacact taaaatcagc 300
agagtggagg ctgaagattt gggagtttat tactgcttgc aaagtacata ttatccgctc 360
acgttcggtg ctgggaccaa gctggagctg aaacgtacgg tggctgcacc atctgtcttc 420
40 atcttcccg ccatctgatga gcagttgaaa tctggaactg cctctgttgt gtgcctgctg 480
aataacttct atcccagaga ggccaaagta cagtggaagg tggataacgc cctccaatcg 540
ggtaactccc aggagagtgt cacagagcag gacagcaagg acagcaccta cagcctcagc 600
agcaccctga cgctgagcaa agcagactac gagaaacaca aagtctacgc ctgcgaagtc 660
45 acccatcagg gcctgagctc gcccgtcaca aagagcttca acaggggaga gtgttga 717
<210> 36
<211> 324
<212> DNA
50 <213> Mus musculus
<400> 36
gacatcaaga tgaccagtc tccatcttcc atgtatgcat ctctaggaga gagagtcact 60
atcacttgca aggcgagtca ggacattaat aactatttaa gctggttcca gcagaaacca 120
55 gggaaatctc ctaagaccct gatctatcgt gcaaacagat tggtagatgg ggtcccatca 180

aggttcagtg gcagtggatc tgggcaagat tattctctca ccatcagcag cctggagtat 240
 gaagatatgg gaattaatta ttgtctacag tgtgatgagt ttctccgtg gacgttcggg 300
 5 ggaggcacca agctggaaat caaa 324
 <210> 37
 <211> 336
 <212> DNA
 10 <213> Mus musculus
 <400> 37
 gatgttgtga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
 atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120
 15 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtctc caaccgattt 180
 tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 agcagagtgg aggctgagga tctgggaggt tatttctgct ctcaaagtac acatgttcgg 300
 tggacgttcg gtggaggcac caagctggaa atcaaa 336
 20 <210> 38
 <211> 705
 <212> DNA
 <213> Mus musculus
 25 <400> 38
 atgagaccct ccattcagtt cctggggctc ttgttgttct ggcttcatgg tgttcagtgt 60
 gacatccaga tgacacagtc tccatcctca ctgtctgcat ctctgggagg caaagtcacc 120
 atcacttgca aggcaagtca ggacattaac aagaatatag tttggtacca acacaagcct 180
 30 ggaaaaggtc ctaggctgct catatggtac acatctacat tacagccagg catcccatca 240
 aggttcagtg gaagtgggtc tgggagagat tattccttca gcatcagcaa cctggagcct 300
 gaagatatgg caacttatta ctgtctacag tatgataatc ttccacggac gttcgggtgga 360
 ggcaccaaac tggaaatcaa acgtacggtg gctgcaccat ctgtcttcat ctcccgcca 420
 35 tctgatgagc agttgaaatc tggaaactgcc tctgttgtgt gcctgctgaa taacttctat 480
 cccagagagg ccaaagtaca gtggaagggtg gataacgcc tccaatcggg taactccag 540
 gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg 600
 40 ctgagcaaag cagactacga gaaacacaaa gtctacgcct gcgaagtcac ccatcagggc 660
 ctgagctcgc ccgtcacaaa gagcttcaac aggggagagt gttga 705
 <210> 39
 <211> 321
 <212> DNA
 45 <213> Mus musculus
 <400> 39
 gacatccaga tgacacagtc tccatcctca ctgtctgcat ctctgggagg caaagtcacc 60
 atcacttgca aggcaagtca ggacattaac aagaatataa tttggtacca acacaagcct 120
 50 ggaaaaggtc ctaggctgct catatggtac acatctacat tacagccagg catcccatca 180
 aggttcagtg gaagtgggtc tgggagagat tattccttca gcatcagcaa cctggagcct 240
 gaagatatgg caacttatta ctgtctacag tatgataatc ttccacggac gttcgggtgga 300
 55 ggcaccaagc tggaaatcaa a 321

<210> 40

<211> 720

<212> DNA

<213> Mus musculus

<400> 40

atgaggttct ctgctcagct tctggggctg cttgtgctct ggatccctgg atccactgca 60
 gatattgtga tgacgcaggc tgcattctcc aatccagtca ctcttggaac atcaacttcc 120
 atctcctgca ggtctagtaa gagtctccta catagtaatg gcatcactta tttgtattgg 180
 tatctgcaga agccaggcca gtctcctcag ctcttgattt atcagatgtc caaccttgcc 240
 tcaggagtcc cagacagggt cagtagcagt gggtcaggaa ctgatttcac actgagaatc 300
 agcagagtgg aggctgagga tgtgggtgtt tattactgtg ctcaaaatct agaacttccg 360
 tatacgttcg gatcggggac caagctggaa ataaaacgta cgggtggctgc accatctgtc 420
 ttcattctcc cgccatctga tgagcagttg aaatctggaa ctgcctctgt tgtgtgcctg 480
 ctgaataact tctatcccag agaggccaaa gtacagtggg aggtggataa cgccctccaa 540
 tcgggtaact cccaggagag tgtcacagag caggacagca aggacagcac ctacagcctc 600
 agcagcaccg tgacgctgag caaagcagac tacgagaaac acaaagtcta cgctgcgaa 660
 gtcacccatc agggcctgag ctgcgccgtc acaaagagct tcaacagggg agagtgttga 720

<210> 41

<211> 336

<212> DNA

<213> Mus musculus

<400> 41

gatattgtga tgacgcaggc tgcattctcc aatccagtca ctcttggaac atcagcttcc 60
 atctcctgca ggtctagtaa gagtctccta catagtaatg gcatcactta tttgtattgg 120
 tttctgcaga agccaggcca gtctcctcag ctcttgattt atcagatgtc caaccttgcc 180
 tcaggagtcc cagacagggt cagtagcagt gggtcaggaa ctgatttcac actgagaatc 240
 agcagagtgg aggctgagga tgtgggtgtt tattactgtg ctcaaaatct agaacttccg 300
 tatacgttcg gatcggggac caagctggaa ataaaa 336

<210> 42

<211> 321

<212> DNA

<213> Mus musculus

<400> 42

gatattgtgc taactcagtc tccagccacc ctgtctgtga ctccaggaga cagagtcagt 60
 ctttctcctgca gggccagcca tagtattagc aacttcctac actggtatcc acaaaaatca 120
 catgagtctc caaggcttct catcaagtat gcttccagtc ccatctctgg gatcccctcc 180
 aggttcagtg gcaatggatc agggacagat ttcaactctca gtatcaacag tgtggagact 240
 gaagattttg gaatgtattt ctgtcaacag agtaacatct ggctcgctcac gttcgggtgct 300
 gggaccaagc tggagctgaa a 321

<210> 43

<211> 333

<212> DNA

<213> Mus musculus

<400> 43

5 gacattgtgc tcaccaatc tccaacttct ttggctgtgt ctctagggca gagtgtcacc 60
 atctcctgca gagccagtga aagtgttgaa tattatggca ctagtttaat gcagtgggtac 120
 caacagaaac caggacagcc acccaaactc ctcatctatg gtgcatccaa cgtagaatct 180
 ggggtccctg ccaggtttag tggcagtggg tctgggacag acttcagcct caacatccat 240
 10 cctgtggagg aggatgatat tgcaatgtat ttctgtcagc aaagtaggaa ggttccgtat 300
 acgttcggat cggggaccaa gctggaaata aaa 333

<210> 44

<211> 238

15 <212> PRT

<213> Mus musculus

<400> 44

Met Ser Pro Ala Gln Phe Leu Phe Leu Leu Val Leu Trp Ile Arg Glu
 20 1 5 10 15
 Thr Asn Gly Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val
 20 25 30
 Thr Ile Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu
 25 35 40 45
 Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro
 50 55 60
 Gly Gln Ser Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser
 30 65 70 75 80
 Gly Ala Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr
 85 90 95
 Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Ile Tyr Tyr Cys
 35 100 105 110
 Trp Gln Gly Thr His Phe Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu
 115 120 125
 Glu Leu Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro
 40 130 135 140
 Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu
 145 150 155 160
 Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn
 45 165 170 175
 Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser
 180 185 190
 Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala
 50 195 200 205
 Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly
 210 215 220
 Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 55

EP 1 674 111 A1

225 230 235

<210> 45

<211> 112

5 <212> PRT

<213> Mus musculus

<400> 45

10 Asp Val Val Met Thr Gln Ser Pro Leu Thr Leu Ser Ile Thr Ile Gly

1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser

20 25 30

15 Asp Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser

35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro

50 55 60

20 Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Ser Leu Lys Ile

65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Ile Tyr Tyr Cys Trp Gln Gly

85 90 95

25 Thr His Phe Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys

100 105 110

<210> 46

<211> 238

30 <212> PRT

<213> Mus musculus

<400> 46

35 Met Ser Pro Val Gln Phe Leu Phe Leu Leu Met Leu Trp Ile Gln Glu

1 5 10 15

Thr Asn Gly Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Ser Val

20 25 30

40 Thr Ile Gly Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu

35 40 45

Leu Tyr Ser Asn Gly Lys Thr Tyr Leu Asn Trp Leu Gln Gln Arg Pro

50 55 60

45 Gly Gln Ala Pro Lys His Leu Met Tyr Gln Val Ser Lys Leu Asp Pro

65 70 75 80

Gly Ile Pro Asp Arg Phe Ser Gly Ser Gly Ser Glu Thr Asp Phe Thr

85 90 95

50 Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys

100 105 110

Leu Gln Ser Thr Tyr Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu

115 120 125

55 Glu Leu Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro

EP 1 674 111 A1

130 135 140
 Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu
 145 150 155 160
 5 Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn
 165 170 175
 Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser
 10 180 185 190
 Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala
 195 200 205
 Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly
 15 210 215 220
 Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235
 <210> 47
 20 <211> 108
 <212> PRT
 <213> Mus musculus
 <400> 47
 25 Asp Ile Lys Met Thr Gln Ser Pro Ser Ser Met Tyr Ala Ser Leu Gly
 1 5 10 15
 Glu Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile Asn Asn Tyr
 20 25 30
 30 Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ser Pro Lys Thr Leu Ile
 35 40 45
 Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 35 Ser Gly Ser Gly Gln Asp Tyr Ser Leu Thr Ile Ser Ser Leu Glu Tyr
 65 70 75 80
 Glu Asp Met Gly Ile Asn Tyr Cys Leu Gln Cys Asp Glu Phe Pro Pro
 85 90 95
 40 Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105
 <210> 48
 45 <211> 112
 <212> PRT
 <213> Mus musculus
 <400> 48
 50 Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15
 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 55 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser

EP 1 674 111 A1

35 40 45
 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 5 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
 10 85 90 95
 Thr His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 49
 15 <211> 234
 <212> PRT
 <213> Mus musculus
 <400> 49
 20 Met Arg Pro Ser Ile Gln Phe Leu Gly Leu Leu Leu Phe Trp Leu His
 1 5 10 15
 Gly Val Gln Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 20 25 30
 25 Ala Ser Leu Gly Gly Lys Val Thr Ile Thr Cys Lys Ala Ser Gln Asp
 35 40 45
 Ile Asn Lys Asn Ile Val Trp Tyr Gln His Lys Pro Gly Lys Gly Pro
 50 55 60
 30 Arg Leu Leu Ile Trp Tyr Thr Ser Thr Leu Gln Pro Gly Ile Pro Ser
 65 70 75 80
 Arg Phe Ser Gly Ser Gly Ser Gly Arg Asp Tyr Ser Phe Ser Ile Ser
 85 90 95
 35 Asn Leu Glu Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp
 100 105 110
 Asn Leu Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
 115 120 125
 40 Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 130 135 140
 Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 145 150 155 160
 45 Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 165 170 175
 Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 180 185 190
 50 Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 195 200 205
 His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 55 210 215 220

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys

225 230

<210> 50

<211> 107

<212> PRT

<213> Mus musculus

<400> 50

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Leu Gly

1 5 10 15

Gly Lys Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile Asn Lys Asn

20 25 30

Ile Ile Trp Tyr Gln His Lys Pro Gly Lys Gly Pro Arg Leu Leu Ile

35 40 45

Trp Tyr Thr Ser Thr Leu Gln Pro Gly Ile Pro Ser Arg Phe Ser Gly

50 55 60

Ser Gly Ser Gly Arg Asp Tyr Ser Phe Ser Ile Ser Asn Leu Glu Pro

65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Leu Pro Arg

85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys

100 105

<210> 51

<211> 239

<212> PRT

<213> Mus musculus

<400> 51

Met Arg Phe Ser Ala Gln Leu Leu Gly Leu Leu Val Leu Trp Ile Pro

1 5 10 15

Gly Ser Thr Ala Asp Ile Val Met Thr Gln Ala Ala Phe Ser Asn Pro

20 25 30

Val Thr Leu Gly Thr Ser Thr Ser Ile Ser Cys Arg Ser Ser Lys Ser

35 40 45

Leu Leu His Ser Asn Gly Ile Thr Tyr Leu Tyr Trp Tyr Leu Gln Lys

50 55 60

Pro Gly Gln Ser Pro Gln Leu Leu Ile Tyr Gln Met Ser Asn Leu Ala

65 70 75 80

Ser Gly Val Pro Asp Arg Phe Ser Ser Ser Gly Ser Gly Thr Asp Phe

85 90 95

Thr Leu Arg Ile Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr

100 105 110

Cys Ala Gln Asn Leu Glu Leu Pro Tyr Thr Phe Gly Ser Gly Thr Lys

115 120 125

EP 1 674 111 A1

Leu Glu Ile Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro
 130 135 140
 5 Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu
 145 150 155 160
 Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp
 165 170 175
 10 Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp
 180 185 190
 Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys
 195 200 205
 15 Ala Asp Tyr Glu Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln
 210 215 220
 Gly Leu Ser Ser Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230 235
 20 <210> 52
 <211> 112
 <212> PRT
 <213> Mus musculus
 25 <400> 52
 Asp Ile Val Met Thr Gln Ala Ala Phe Ser Asn Pro Val Thr Leu Gly
 1 5 10 15
 Thr Ser Ala Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu His Ser
 30 20 25 30
 Asn Gly Ile Thr Tyr Leu Tyr Trp Phe Leu Gln Lys Pro Gly Gln Ser
 35 35 40 45
 Pro Gln Leu Leu Ile Tyr Gln Met Ser Asn Leu Ala Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Ser Ser Gly Ser Gly Thr Asp Phe Thr Leu Arg Ile
 65 70 75 80
 40 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ala Gln Asn
 85 90 95
 Leu Glu Leu Pro Tyr Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 45 <210> 53
 <211> 107
 <212> PRT
 <213> Mus musculus
 50 <400> 53
 Asp Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
 1 5 10 15
 Asp Arg Val Ser Leu Ser Cys Arg Ala Ser His Ser Ile Ser Asn Phe
 55 20 25 30

Leu His Trp Tyr Pro Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
 35 40 45
 5 Lys Tyr Ala Ser Gln Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60
 Asn Gly Ser Gly Thr Asp Phe Thr Leu Ser Ile Asn Ser Val Glu Thr
 65 70 75 80
 10 Glu Asp Phe Gly Met Tyr Phe Cys Gln Gln Ser Asn Ile Trp Ser Leu
 85 90 95
 Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105
 15 <210> 54
 <211> 111
 <212> PRT
 <213> Mus musculus
 20 <400> 54
 Asp Ile Val Leu Thr Gln Ser Pro Thr Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Gln Ser Val Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Glu Tyr Tyr
 25 20 25 30
 Gly Thr Ser Leu Met Gln Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 30 Lys Leu Leu Ile Tyr Gly Ala Ser Asn Val Glu Ser Gly Val Pro Ala
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Ser Leu Asn Ile His
 65 70 75 80
 35 Pro Val Glu Glu Asp Asp Ile Ala Met Tyr Phe Cys Gln Gln Ser Arg
 85 90 95
 Lys Val Pro Tyr Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 40 <210> 55
 <211> 333
 <212> DNA
 <213> Mus musculus
 45 <400> 55
 cagatccagt tggagcagtc tggacctgag ctgaagaagc ctggagagac agtcaagatc 60
 tcttgcaagg cttctggtta tattttcaga gactattcaa tgcactgggt gaagcaggct 120
 ccaggaaagg gtttaaagtg gatgggctgg ataaacactg agacgggtga gccaacatat 180
 50 gcagatgact tcaagggacg gtttgccctt tctttggaaa cctctgccag cactgcctat 240
 ttgcagatca acaacctcaa aaatgaggac acggctacat atttctgtac tagcctttac 300
 tggggccaag ggactctggt cactgtctct gca 333
 <210> 56
 55 <211> 372

<212> DNA

<213> Mus musculus

<400> 56

5 cagggtcactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
 acttgttctt tctctgggtt ttcactgagc acttatggta tgggtgtagg ttggattcgt 120
 cagccttcag ggaagggtct ggagtggctg gccaacattt ggtggcatga tgataagtac 180
 10 tataactcag ccctgaagag ccggctcaca atctccaagg atatctcaa caaccaggta 240
 ttcctcaaga tctccagtgt ggacactgca gatactgcca catactactg tgctcaaata 300
 gccctcgat ataataagta cgaaggcttt tttgctttct ggggccaagg gactctggtc 360
 actgtctctg ca 372

<210> 57

<211> 345

<212> DNA

<213> Mus musculus

<400> 57

20 cagggttcaac tgcagcagtc tggggctgag ctgggtgaggc ctggggcttc agtgaagctg 60
 tcctgcaagg cttcgggcta cacatttact gactatgaaa tgcactgggt gaagcagaca 120
 cctgtgcatg gcctaaaatg gattggagct cttgatccta aaactgggtga tactgcctac 180
 25 agtcagaagt tcaagggcaa ggccacactg actgcagaca aatcctccag cacagcctac 240
 atggagctcc gcagcctgac atctgaggac tctgccgtct attactgtac aagattctac 300
 tcctatactt actggggcca agggactctg gtcactgtct ctgca 345

<210> 58

<211> 357

<212> DNA

<213> Mus musculus

<400> 58

35 gaggtgcagc ttgttgagac tgggtggagga ctgggtgcagc ctgaagggtc attgaaactc 60
 tcatgtgcag cttctggatt cagcttcaat atcaatgcca tgaactgggt ccgccaggct 120
 ccaggaaagg gtttggaatg ggttgctcgc ataagaagtg aaagtaataa ttatgcaaca 180
 40 tattatggcg attcagtgaag agacagggttc accatctcca gagatgattc acaaaacatg 240
 ctctatctac aaatgaacaa cttgaaaact gaggacacag ccatatatta ctgtgtgaga 300
 gaggttaacta catcgtttgc ttattggggc caagggactc tggtcactgt ctctgca 357

<210> 59

<211> 369

<212> DNA

<213> Mus musculus

<400> 59

50 gaggtgcagc ttgttgagac tgggtggagga ttgggtgcagc ctaaagggtc attgaaactc 60
 tcatgtgcag cctctggatt caccttcaat gccagtcca tgaactgggt ccgccaggct 120
 ccaggaaagg gtttggaatg ggttgctcgc ataagaagta aaagtaataa ttatgcaata 180
 tattatgccg attcagtgaag agacagggttc accatctcca gagatgattc acaagcatg 240
 55 ctctatctgc aaatgaacaa cttgaaaact gaggacacag ccatgtatta ctgtgtgaga 300

gatccgggct actatggtaa cccctgggtt gcttactggg gccaaaggac tctgggtcact 360
gtctctgca 369

5

<210> 60

<211> 111

<212> PRT

<213> Mus musculus

10

<400> 60

Gln Ile Gln Leu Glu Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu

1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ile Phe Arg Asp Tyr

15

20 25 30

Ser Met His Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Met

35 40 45

Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe

20

50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr

65 70 75 80

Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys

25

85 90 95

Thr Ser Leu Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala

100 105 110

30

<210> 61

<211> 124

<212> PRT

<213> Mus musculus

35

<400> 61

Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln

1 5 10 15

Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Tyr

40

20 25 30

Gly Met Gly Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu

35 40 45

Trp Leu Ala Asn Ile Trp Trp His Asp Asp Lys Tyr Tyr Asn Ser Ala

45

50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Ile Ser Asn Asn Gln Val

65 70 75 80

Phe Leu Lys Ile Ser Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr

50

85 90 95

Cys Ala Gln Ile Ala Pro Arg Tyr Asn Lys Tyr Glu Gly Phe Phe Ala

100 105 110

Phe Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala

55

115 120

<210> 62

<211> 115

<212> PRT

<213> Mus musculus

<400> 62

Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly Ala
 1 5 10 15
 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Glu Met His Trp Val Lys Gln Thr Pro Val His Gly Leu Lys Trp Ile
 35 40 45
 Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115

<210> 63

<211> 119

<212> PRT

<213> Mus musculus

<400> 63

Glu Val Gln Leu Val Glu Thr Gly Gly Gly Leu Val Gln Pro Glu Gly
 1 5 10 15
 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Ser Phe Asn Ile Asn
 20 25 30
 Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ala Arg Ile Arg Ser Glu Ser Asn Asn Tyr Ala Thr Tyr Tyr Gly Asp
 50 55 60
 Ser Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Gln Asn Met
 65 70 75 80
 Leu Tyr Leu Gln Met Asn Asn Leu Lys Thr Glu Asp Thr Ala Ile Tyr
 85 90 95
 Tyr Cys Val Arg Glu Val Thr Thr Ser Phe Ala Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ala
 115

<210> 64

<211> 123

<212> PRT

<213> Mus musculus

<400> 64

Glu Val Gln Leu Val Glu Thr Gly Gly Gly Leu Val Gln Pro Lys Gly

1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Asn Ala Ser

20 25 30

Ala Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val

35 40 45

Ala Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Ile Tyr Tyr Ala Asp

50 55 60

Ser Val Lys Asp Arg Phe Thr Ile Ser Arg Asp Asp Ser Gln Ser Met

65 70 75 80

Leu Tyr Leu Gln Met Asn Asn Leu Lys Thr Glu Asp Thr Ala Met Tyr

85 90 95

Tyr Cys Val Arg Asp Pro Gly Tyr Tyr Gly Asn Pro Trp Phe Ala Tyr

100 105 110

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala

115 120

<210> 65

<211> 336

<212> DNA

<213> Mus musculus

<400> 65

gatgttgtga tgaccagac tccactcact ttgtcgggta cccttggaca accagcctcc 60

atctcttgca agtcaagtca gagcctotta catagtgatg gaaagacatt ttgaattgg 120

ttattacaga ggccaggcca gtctccaaag cgcctaattct atctgggtgct tagactggac 180

tctggagtcc ctgacagggt cactggcagt ggatcagggg cagatttcac actgaaaatc 240

agcagagtgg aggctgagga tttgggagtt tattattgct gccaagggtac acattttcct 300

cggacgttcg gtggaggcac caggctggaa atcaaa 336

<210> 66

<211> 336

<212> DNA

<213> Mus musculus

<400> 66

gatgttttga tgaccocaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60

atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120

tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180

tctgggggtcc cagacagggt cagtggcagt ggatcagggg cagatttcac actcaagatc 240

agcagagtgg aggctgagga tctgggagtt tattactgct ttcaagggtc acatgttccg 300

tggacgttcg gtggaggcac caagctggaa atcaaa 336
 <210> 67
 <211> 336
 <212> DNA
 <213> Mus musculus
 <400> 67
 10 gatgttgtga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
 atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120
 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180
 tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 15 agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaaatac acatgttcct 300
 cctacgttcg gatcggggac caagctggaa ataaaa 336
 <210> 68
 <211> 336
 20 <212> DNA
 <213> Mus musculus
 <400> 68
 25 gatattgtga tgactcagtc tgcaccctct gtacctgtca ctcttggaga gtcagtatcc 60
 atctcctgca agtctagtaa gagtctcctg catagtaatg gcaacactta cttgaattgg 120
 ttcctgcaga ggccaggcca gtctcctcaa ctctgattt attggatgtc caaccttgcc 180
 tcaggagtcc cagacagggt cagtggcagt gggtcaggaa ctgctttcac actgagaatc 240
 30 agtagagtgg aggctgagga tgtgggtgtt tattactgta tgcaacatat agaataccct 300
 ttcacgttcg gcacggggac aaaattggaa ataaaa 336
 <210> 69
 <211> 336
 35 <212> DNA
 <213> Mus musculus
 <400> 69
 gatattgtga tgacgcaggc tgcattctcc aatccagtca ctcttggaa atcagcttcc 60
 40 atctcctgca ggtctagtaa gagtctccta catagttagt acatcactta tttgtattgg 120
 tatctgcaga agccaggcca gtctcctcag ctctgattt atcagatgtc caaccttgcc 180
 tcaggagtcc cagacagggt cagtagcagt gggtcaggaa ctgatttcac actgagaatc 240
 agcagagtgg aggctgagga tgtgggtgtt tattactgtg ctcaaaatct agaacttct 300
 45 ccgacgttcg gtggaggcac caagctggaa atcaaa 336
 <210> 70
 <211> 318
 <212> DNA
 50 <213> Mus musculus
 <400> 70
 caaattgttc tcaccagtc tccagcaatc atgtotgeat ttccagggga gaaggtcacc 60
 55 atgacctgca gtgccagctc aagtgttagt tacatgtact ggtaccagca gaagtcagga 120
 tcctccccca gactcctgat ttatgacaca tccaacctgg cttctggagt ccctgttcgc 180

EP 1 674 111 A1

ttcagtggca gtgggtctgg gacctcttac tctctcacia tcagccgaat ggaggctgaa 240
gatgctgcca cttattactg ccagcagtgg agtagttacc cgctcacgtt cgggtgggtggg 300
accgagctgg agctgaaa 318

5
<210> 71
<211> 112
<212> PRT
10
<213> Mus musculus
<400> 71
Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Leu Gly
1 5 10 15
15 Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu His Ser
20 25 30
Asp Gly Lys Thr Phe Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
35 40 45
20 Pro Lys Arg Leu Ile Tyr Leu Val Ser Arg Leu Asp Ser Gly Val Pro
50 55 60
Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80
25 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Cys Gln Gly
85 90 95
Thr His Phe Pro Arg Thr Phe Gly Gly Gly Thr Arg Leu Glu Ile Lys
100 105 110
30
<210> 72
<211> 112
<212> PRT
35
<213> Mus musculus
<400> 72
Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15
40 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
20 25 30
Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45
45 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80
50 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly
85 90 95
Ser His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110
55
<210> 73

<211> 112

<212> PRT

<213> Mus musculus

<400> 73

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly

1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser

20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser

35 40 45

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro

50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile

65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Asn

85 90 95

Thr His Val Pro Pro Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys

100 105 110

<210> 74

<211> 112

<212> PRT

<213> Mus musculus

<400> 74

Asp Ile Val Met Thr Gln Ser Ala Pro Ser Val Pro Val Thr Pro Gly

1 5 10 15

Glu Ser Val Ser Ile Ser Cys Lys Ser Ser Lys Ser Leu Leu His Ser

20 25 30

Asn Gly Asn Thr Tyr Leu Asn Trp Phe Leu Gln Arg Pro Gly Gln Ser

35 40 45

Pro Gln Leu Leu Ile Tyr Trp Met Ser Asn Leu Ala Ser Gly Val Pro

50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Ala Phe Thr Leu Arg Ile

65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln His

85 90 95

Ile Glu Tyr Pro Phe Thr Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys

100 105 110

<210> 75

<211> 112

<212> PRT

<213> Mus musculus

<400> 75

Asp Ile Val Met Thr Gln Ala Ala Phe Ser Asn Pro Val Thr Leu Gly
 1 5 10 15
 5 Thr Ser Ala Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu His Ser
 20 25 30
 Tyr Asp Ile Thr Tyr Leu Tyr Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 10 Pro Gln Leu Leu Ile Tyr Gln Met Ser Asn Leu Ala Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Ser Ser Gly Ser Gly Thr Asp Phe Thr Leu Arg Ile
 65 70 75 80
 15 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ala Gln Asn
 85 90 95
 Leu Glu Leu Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 20 <210> 76
 <211> 106
 <212> PRT
 <213> Mus musculus
 25 <400> 76
 Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Phe Pro Gly
 1 5 10 15
 30 Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
 20 25 30
 Tyr Trp Tyr Gln Gln Lys Ser Gly Ser Ser Pro Arg Leu Leu Ile Tyr
 35 40 45
 35 Asp Thr Ser Asn Leu Ala Ser Gly Val Pro Val Arg Phe Ser Gly Ser
 50 55 60
 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Arg Met Glu Ala Glu
 65 70 75 80
 40 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Tyr Pro Leu Thr
 85 90 95
 Phe Gly Gly Gly Thr Glu Leu Glu Leu Lys
 100 105
 45 <210> 77
 <211> 345
 <212> DNA
 <213> Artificial Sequence
 50 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 77
 caggtgcagc tggaggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 55 tctctcaagg cttctggata caccttcacc gactatgaaa tgcaactgggt gcgacaggcc 120

cctggacaag ggcttgagtg gatgggagct cttgataccta aaactgggtga tactgcctac 180
 agtcagaagt tcaagggcag agtcacgatt accgcggacg aatccacgag cacagcctac 240
 5 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagattctac 300
 tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345
 <210> 78
 <211> 345
 10 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Mouse-human chimeric antibody H chain
 15 <400> 78
 caggtgcagc tgggtggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagct cttgataccta aaactgggtga tactgcctac 180
 20 agtcagaagt tcaagggcag agtcacgctg accgcggacg aatccacgag cacagcctac 240
 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac 300
 tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345
 <210> 79
 25 <211> 345
 <212> DNA
 <213> Artificial Sequence
 <220>
 30 <223> Mouse-human chimeric antibody H chain
 <400> 79
 caggtgcagc tgggtggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120
 35 cctggacaag ggcttgagtg gatgggagct cttgataccta aaactgggtga tactgcctac 180
 agtcagaagt tcaagggcag agtcacgctg accgcggaca aatccacgag cacagcctac 240
 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac 300
 40 tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345
 <210> 80
 <211> 345
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 80
 50 caggtgcagc tgggtggagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagct cttgataccta aaactgggtga tactgcctac 180
 agtcagaagt tcaagggcag agtcacgctg accgcggaca aatccacgag cacagcctac 240
 55 atggagctga gcagcctgac atctgaggac acggccgtgt attactgtac aagattctac 300

tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345

<210> 81

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 81

caggtgcagc tgggtgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60

tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120

cctggacaag ggcttgagtg gatgggagct cttgatccta aaactgggtga tactgcctac 180

agtcagaagt tcaagggcag agtcacgctg accgcgagc aatccacgag cacagcctac 240

atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac 300

tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345

<210> 82

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 82

caggtgcagc tgggtgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60

tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120

cctggacaag ggcttgagtg gatgggagct cttgatccta aaactgggtga tactgcctac 180

agtcagaagt tcaagggcag agtcacgctg accgcgagc aatccacgag cacagcctac 240

atggagctga gcagcctgag atctgaggac acggccgtgt attactgtac aagattctac 300

tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345

<210> 83

<211> 345

<212> DNA

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 83

caggtgcagc tgggtgcagtc tggagctgag gtgaagaagc ctggggcctc agtgaaggtc 60

tcctgcaagg cttctggata caccttcacc gactatgaaa tgcactgggt gcgacaggcc 120

cctggacaag ggcttgagtg gatgggagct cttgatccta aaactgggtga tactgcctac 180

agtcagaagt tcaagggcag agtcacgctg accgcgagc aatccacgag cacagcctac 240

atggagctga gcagcctgac atctgaggac acggccgtgt attactgtac aagattctac 300

tcctatactt actggggcca gggaaccctg gtcaccgtct cctca 345

<210> 84

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 84

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 Lys Gly Arg Val Thr Ile Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 25 100 105 110
 Val Ser Ser
 115

<210> 85

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 85

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 Lys Gly Arg Val Thr Leu Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
 50 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 55 100 105 110

Val Ser Ser

115

<210> 86

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 86

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr

20 25 30

Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met

35 40 45

Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe

50 55 60

Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys

85 90 95

Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr

100 105 110

Val Ser Ser

115

<210> 87

<211> 115

<212> PRT

<213> Artificial Sequence

<220>

<223> Mouse-human chimeric antibody H chain

<400> 87

Gln Val Gln Leu Val Glu Ser Gly Ala Glu Val Lys Lys Pro Gly Ala

1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr

20 25 30

Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met

35 40 45

Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe

50 55 60

Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr

65 70 75 80

Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 5 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ser
 115
 10 <210> 88
 <211> 115
 <212> PRT
 <213> Artificial Sequence
 15 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 88
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 20 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 25 Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 30 Lys Gly Arg Val Thr Leu Thr Ala Asp Glu Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 35 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ser
 115
 40 <210> 89
 <211> 115
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 89
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 50 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 55 Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

EP 1 674 111 A1

Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 5 Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 10 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ser
 115
 15 <210> 90
 <211> 115
 <212> PRT
 <213> Artificial Sequence
 20 <220>
 <223> Mouse-human chimeric antibody H chain
 <400> 90
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 25 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 Glu Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 30 35 40 45
 Gly Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe
 50 55 60
 35 Lys Gly Arg Val Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 40 Thr Arg Phe Tyr Ser Tyr Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ser
 115
 45 <210> 91
 <211> 336
 <212> DNA
 <213> Artificial Sequence
 50 <220>
 <223> Mouse-human chimeric antibody L chain
 <400> 91
 gatgttgatga tgactcagtc tccactctcc ctgcccgtca cccctggaga gccggcctcc 60
 55 atctcctgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120

tacctgcaga agccagggca gtctccacag ctctgatct ataaagtttc caaccgattt 180
 tctggggtcc ctgacagggt cagtggcagt ggatcaggca cagattttac actgaaaatc 240
 5 agcagagtgg aggctgagga tgttgggggt tattactgct ctcaaaatac acatgttcct 300
 cctacgtttg gccaggggac caagctggag atcaaa 336
 <210> 92
 <211> 112
 10 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> Mouse-human chimeric antibody L chain
 15 <400> 92
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 20 25 30
 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 25 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 30 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 35 <210> 93
 <211> 14
 <212> PRT
 <213> homo sapiens
 40 <400> 93
 Gly Asn Ser Gln Gln Ala Thr Pro Lys Asp Asn Glu Ile Ser
 1 5 10
 45 <210> 94
 <211> 8
 <212> PRT
 <213> homo sapiens
 <400> 94
 50 Gly Asn Ser Gln Gln Ala Thr Pro
 1 5
 <210> 95
 <211> 8
 55 <212> PRT

<213> homo sapiens
 <400> 95
 5 Gln Gln Ala Thr Pro Lys Asp Asn
 1 5
 <210> 96
 <211> 8
 10 <212> PRT
 <213> homo sapiens
 <400> 96
 Thr Pro Lys Asp Asn Glu Ile Ser
 15 1 5
 <210> 97
 <211> 10
 <212> PRT
 20 <213> homo sapiens
 <400> 97
 Ala Thr Pro Lys Asp Asn Glu Ile Ser Thr
 1 5 10
 25 <210> 98
 <211> 10
 <212> PRT
 <213> homo sapiens
 30 <400> 98
 Pro Lys Asp Asn Glu Ile Ser Thr Phe His
 1 5 10
 35 <210> 99
 <211> 10
 <212> PRT
 <213> homo sapiens
 <400> 99
 40 Asp Asn Glu Ile Ser Thr Phe His Asn Leu
 1 5 10
 <210> 100
 <211> 10
 45 <212> PRT
 <213> homo sapiens
 <400> 100
 50 Glu Ile Ser Thr Phe His Asn Leu Gly Asn
 1 5 10
 <210> 101
 <211> 27
 55 <212> PRT

<213> homo sapiens

<400> 101

Gly Asn Ser Gln Gln Ala Thr Pro Lys Asp Asn Glu Ile Ser Thr Phe

1 5 10 15

His Asn Leu Gly Asn Val His Ser Pro Leu Lys

20 25

<210> 102

<211> 14

<212> PRT

<213> homo sapiens

<400> 102

Ser Thr Phe His Asn Leu Gly Asn Val His Ser Pro Leu Lys

1 5 10

<210> 103

<211> 5

<212> PRT

<213> Mus musculus

<400> 103

Asn Tyr Ala Met Ser

1 5

<210> 104

<211> 17

<212> PRT

<213> Mus musculus

<400> 104

Ala Ile Asn Asn Asn Gly Asp Asp Thr Tyr Tyr Leu Asp Thr Val Lys

1 5 10 15

Asp

<210> 105

<211> 5

<212> PRT

<213> Mus musculus

<400> 105

Gln Gly Gly Ala Tyr

1 5

<210> 106

<211> 7

<212> PRT

<213> Mus musculus

<400> 106

Thr Tyr Gly Met Gly Val Gly

1 5
 <210> 107
 5 <211> 16
 <212> PRT
 <213> Mus musculus
 <400> 107
 10 Asn Ile Trp Trp Tyr Asp Ala Lys Tyr Tyr Asn Ser Asp Leu Lys Ser
 1 5 10 15
 <210> 108
 <211> 8
 15 <212> PRT
 <213> Mus musculus
 <400> 108
 Met Gly Leu Ala Trp Phe Ala Tyr
 20 1 5
 <210> 109
 <211> 7
 <212> PRT
 25 <213> Mus musculus
 <400> 109
 Ile Tyr Gly Met Gly Val Gly
 30 1 5
 <210> 110
 <211> 16
 <212> PRT
 35 <213> Mus musculus
 <400> 110
 Asn Ile Trp Trp Asn Asp Asp Lys Tyr Tyr Asn Ser Ala Leu Lys Ser
 1 5 10 15
 <210> 111
 40 <211> 8
 <212> PRT
 <213> Mus musculus
 <400> 111
 45 Ile Gly Tyr Phe Tyr Phe Asp Tyr
 1 5
 <210> 112
 <211> 5
 50 <212> PRT
 <213> Mus musculus
 <400> 112
 55 Gly Tyr Trp Met His

1 5
 <210> 113
 5 <211> 17
 <212> PRT
 <213> Mus musculus
 <400> 113
 10 Ala Ile Tyr Pro Gly Asn Ser Asp Thr Asn Tyr Asn Gln Lys Phe Lys
 1 5 10 15
 Gly
 15 <210> 114
 <211> 10
 <212> PRT
 <213> Mus musculus
 20 <400> 114
 Ser Gly Asp Leu Thr Gly Gly Leu Ala Tyr
 1 5 10
 25 <210> 115
 <211> 5
 <212> PRT
 <213> Mus musculus
 30 <400> 115
 Ser Tyr Ala Met Ser
 1 5
 <210> 116
 35 <211> 17
 <212> PRT
 <213> Mus musculus
 <400> 116
 40 Ala Ile Asn Ser Asn Gly Gly Thr Thr Tyr Tyr Pro Asp Thr Met Lys
 1 5 10 15
 Asp
 45 <210> 117
 <211> 13
 <212> PRT
 <213> Mus musculus
 50 <400> 117
 His Asn Gly Gly Tyr Glu Asn Tyr Gly Trp Phe Ala Tyr
 1 5 10
 55 <210> 118
 <211> 5

<212> PRT
 <213> Mus musculus
 <400> 118
 5 Ser Tyr Trp Met His
 1 5
 <210> 119
 <211> 17
 10 <212> PRT
 <213> Mus musculus
 <400> 119
 15 Glu Ile Asp Pro Ser Asp Ser Tyr Thr Tyr Tyr Asn Gln Lys Phe Arg
 1 5 10 15
 Gly
 <210> 120
 <211> 15
 <212> PRT
 <213> Mus musculus
 25 <400> 120
 Ser Asn Leu Gly Asp Gly His Tyr Arg Phe Pro Ala Phe Pro Tyr
 1 5 10 15
 <210> 121
 <211> 17
 <212> PRT
 <213> Mus musculus
 35 <400> 121
 Thr Ile Asp Pro Ser Asp Ser Glu Thr His Tyr Asn Leu Gln Phe Lys
 1 5 10 15
 Asp
 <210> 122
 <211> 15
 <212> PRT
 45 <213> Mus musculus
 <400> 122
 Gly Ala Phe Tyr Ser Ser Tyr Ser Tyr Trp Ala Trp Phe Ala Tyr
 1 5 10 15
 <210> 123
 <211> 5
 <212> PRT
 <213> Mus musculus
 55 <400> 123

Asp Tyr Glu Met His

1 5

<210> 124

<211> 17

<212> PRT

<213> Mus musculus

<400> 124

Ala Leu Asp Pro Lys Thr Gly Asp Thr Ala Tyr Ser Gln Lys Phe Lys

1 5 10 15

Gly

<210> 125

<211> 6

<212> PRT

<213> Mus musculus

<400> 125

Phe Tyr Ser Tyr Thr Tyr

1 5

<210> 126

<211> 5

<212> PRT

<213> Mus musculus

<400> 126

Ile Asn Ala Met Asn

1 5

<210> 127

<211> 19

<212> PRT

<213> Mus musculus

<400> 127

Arg Ile Arg Ser Glu Ser Asn Asn Tyr Ala Thr Tyr Tyr Gly Asp Ser

1 5 10 15

Val Lys Asp

<210> 128

<211> 8

<212> PRT

<213> Mus musculus

<400> 128

Glu Val Thr Thr Ser Phe Ala Tyr

1 5

<210> 129

<211> 5
 <212> PRT
 <213> Mus musculus
 5 <400> 129
 Ala Ser Ala Met Asn
 1 5
 10 <210> 130
 <211> 19
 <212> PRT
 <213> Mus musculus
 15 <400> 130
 Arg Ile Arg Ser Lys Ser Asn Asn Tyr Ala Ile Tyr Tyr Ala Asp Ser
 1 5 10 15
 Val Lys Asp
 20
 <210> 131
 <211> 12
 <212> PRT
 25 <213> Mus musculus
 <400> 131
 Asp Pro Gly Tyr Tyr Gly Asn Pro Trp Phe Ala Tyr
 1 5 10
 30 <210> 132
 <211> 5
 <212> PRT
 <213> Mus musculus
 35 <400> 132
 Asp Tyr Ser Met His
 1 5
 40 <210> 133
 <211> 17
 <212> PRT
 <213> Mus musculus
 45 <400> 133
 Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe Lys
 1 5 10 15
 Gly
 50
 <210> 134
 <211> 2
 <212> PRT
 55 <213> Mus musculus

<400> 134
 Leu Tyr
 1
 5 <210> 135
 <211> 16
 <212> PRT
 10 <213> Mus musculus
 <400> 135
 Asn Ile Trp Trp His Asp Asp Lys Tyr Tyr Asn Ser Ala Leu Lys Ser
 1 5 10 15
 15 <210> 136
 <211> 14
 <212> PRT
 <213> Mus musculus
 20 <400> 136
 Ile Ala Pro Arg Tyr Asn Lys Tyr Glu Gly Phe Phe Ala Phe
 1 5 10
 25 <210> 137
 <211> 16
 <212> PRT
 <213> Mus musculus
 30 <400> 137
 Lys Ser Ser Gln Ser Leu Leu Asp Ser Asp Gly Lys Thr Tyr Leu Asn
 1 5 10 15
 35 <210> 138
 <211> 7
 <212> PRT
 <213> Mus musculus
 40 <400> 138
 Leu Val Ser Lys Leu Asp Ser
 1 5
 45 <210> 139
 <211> 9
 <212> PRT
 <213> Mus musculus
 50 <400> 139
 Trp Gln Gly Thr His Phe Pro Leu Thr
 1 5
 55 <210> 140
 <211> 11
 <212> PRT
 <213> Mus musculus

<400> 140
 Lys Ala Ser Gln Asp Ile Asn Asn Tyr Leu Ser
 1 5 10
 5 <210> 141
 <211> 7
 <212> PRT
 10 <213> Mus musculus
 <400> 141
 Arg Ala Asn Arg Leu Val Asp
 1 5
 15 <210> 142
 <211> 10
 <212> PRT
 <213> Mus musculus
 20 <400> 142
 Leu Gln Cys Asp Glu Phe Pro Pro Trp Thr
 1 5 10
 25 <210> 143
 <211> 16
 <212> PRT
 <213> Mus musculus
 30 <400> 143
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Asn Thr Tyr Leu His
 1 5 10 15
 35 <210> 144
 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 144
 40 Lys Val Ser Asn Arg Phe Ser
 1 5
 45 <210> 145
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 145
 50 Ser Gln Ser Thr His Val Pro Trp Thr
 1 5
 55 <210> 146
 <211> 16
 <212> PRT
 <213> Mus musculus

<400> 146
 Arg Ser Ser Lys Ser Leu Leu His Ser Asn Gly Ile Thr Tyr Leu Tyr
 1 5 10 15
 <210> 147
 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 147
 Gln Met Ser Asn Leu Ala Ser
 1 5
 <210> 148
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 148
 Ala Gln Asn Leu Glu Leu Pro Tyr Thr
 1 5
 <210> 149
 <211> 11
 <212> PRT
 <213> Mus musculus
 <400> 149
 Lys Ala Ser Gln Asp Ile Asn Lys Asn Ile Ile
 1 5 10
 <210> 150
 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 150
 Tyr Thr Ser Thr Leu Gln Pro
 1 5
 <210> 151
 <211> 6
 <212> PRT
 <213> Mus musculus
 <400> 151
 Leu Gln Tyr Asp Asn Leu
 1 5
 <210> 152
 <211> 11
 <212> PRT
 <213> Mus musculus

5 <400> 152
 Arg Ala Ser His Ser Ile Ser Asn Phe Leu His
 1 5 10
 <210> 153
 <211> 7
 <212> PRT
 10 <213> Mus musculus
 <400> 153
 Tyr Ala Ser Gln Ser Ile Ser
 1 5
 15 <210> 154
 <211> 9
 <212> PRT
 <213> Mus musculus
 20 <400> 154
 Gln Gln Ser Asn Ile Trp Ser Leu Thr
 1 5
 25 <210> 155
 <211> 15
 <212> PRT
 <213> Mus musculus
 30 <400> 155
 Arg Ala Ser Glu Ser Val Glu Tyr Tyr Gly Thr Ser Leu Met Gln
 1 5 10 15
 <210> 156
 35 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 156
 40 Gly Ala Ser Asn Val Glu Ser
 1 5
 <210> 157
 <211> 9
 45 <212> PRT
 <213> Mus musculus
 <400> 157
 Gln Gln Ser Arg Lys Val Pro Tyr Thr
 50 1 5
 <210> 158
 <211> 9
 55 <212> PRT
 <213> Mus musculus

<400> 158
 Ser Gln Asn Thr His Val Pro Pro Thr
 1 5
 <210> 159
 <211> 16
 <212> PRT
 <213> Mus musculus
 <400> 159
 Lys Ser Ser Lys Ser Leu Leu His Ser Asn Gly Asn Thr Tyr Leu Asn
 1 5 10 15
 <210> 160
 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 160
 Trp Met Ser Asn Leu Ala Ser
 1 5
 <210> 161
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 161
 Met Gln His Ile Glu Tyr Pro Phe Thr
 1 5
 <210> 162
 <211> 16
 <212> PRT
 <213> Mus musculus
 <400> 162
 Arg Ser Ser Lys Ser Leu Leu His Ser Tyr Asp Ile Thr Tyr Leu Tyr
 1 5 10 15
 <210> 163
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 163
 Ala Gln Asn Leu Glu Leu Pro Pro Thr
 1 5
 <210> 164
 <211> 10
 <212> PRT
 <213> Mus musculus

<400> 164
 Ser Ala Ser Ser Ser Val Ser Tyr Met Tyr
 1 5 10
 5
 <210> 165
 <211> 7
 <212> PRT
 10
 <213> Mus musculus
 <400> 165
 Asp Thr Ser Asn Leu Ala Ser
 1 5
 15
 <210> 166
 <211> 9
 <212> PRT
 <213> Mus musculus
 20
 <400> 166
 Gln Gln Trp Ser Ser Tyr Pro Leu Thr
 1 5
 25
 <210> 167
 <211> 16
 <212> PRT
 <213> Mus musculus
 30
 <400> 167
 Lys Ser Ser Gln Ser Leu Leu His Ser Asp Gly Lys Thr Phe Leu Asn
 1 5 10 15
 35
 <210> 168
 <211> 7
 <212> PRT
 <213> Mus musculus
 <400> 168
 40
 Leu Val Ser Arg Leu Asp Ser
 1 5
 45
 <210> 169
 <211> 6
 <212> PRT
 <213> Mus musculus
 <400> 169
 50
 Cys Gln Gly Thr His Phe
 1 5
 55
 <210> 170
 <211> 16
 <212> PRT
 <213> Mus musculus

<400> 170
 Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu
 1 5 10 15
 <210> 171
 <211> 9
 <212> PRT
 <213> Mus musculus
 <400> 171
 Phe Gln Gly Ser His Val Pro Trp Thr
 1 5
 <210> 172
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer
 <400> 172
 cttgtacaca gtgacggaaa cacctat 27
 <210> 173
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer
 <400> 173
 ataggtgttt ccgtcactgt gtacaag 27
 <210> 174
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 174
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Ala Asn Thr Tyr Leu His
 1 5 10 15
 <210> 175
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 175

Arg Ser Ser Gln Ser Leu Val His Ser Asn Asp Asn Thr Tyr Leu His
 1 5 10 15
 5 <210> 176
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 10 <220>
 <223> mutant antibody L chain
 <400> 176
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Glu Asn Thr Tyr Leu His
 15 1 5 10 15
 <210> 177
 <211> 16
 <212> PRT
 20 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 177
 25 Arg Ser Ser Gln Ser Leu Val His Ser Asn Phe Asn Thr Tyr Leu His
 1 5 10 15
 <210> 178
 30 <211> 16
 <212> PRT
 <213> Artificial Sequence
 <220>
 35 <223> mutant antibody L chain
 <400> 178
 Arg Ser Ser Gln Ser Leu Val His Ser Asn His Asn Thr Tyr Leu His
 40 1 5 10 15
 <210> 179
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 45 <220>
 <223> mutant antibody L chain
 <400> 179
 50 Arg Ser Ser Gln Ser Leu Val His Ser Asn Asn Asn Thr Tyr Leu His
 1 5 10 15
 <210> 180
 <211> 16
 55 <212> PRT
 <213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 180

Arg Ser Ser Gln Ser Leu Val His Ser Asn Thr Asn Thr Tyr Leu His

1 5 10 15

<210> 181

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 181

Arg Ser Ser Gln Ser Leu Val His Ser Asn Gln Asn Thr Tyr Leu His

1 5 10 15

<210> 182

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 182

Arg Ser Ser Gln Ser Leu Val His Ser Asn Gly Ile Asn Thr Tyr Leu

1 5 10 15

His

<210> 183

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 183

Arg Ser Ser Gln Ser Leu Val His Ser Asn Lys Asn Thr Tyr Leu His

1 5 10 15

<210> 184

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 184

Arg Ser Ser Gln Ser Leu Val His Ser Asn Leu Asn Thr Tyr Leu His

EP 1 674 111 A1

1 5 10 15
 <210> 185
 <211> 16
 5 <212> PRT
 <213> Artificial Sequence
 <220>
 10 <223> mutant antibody L chain
 <400> 185
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Ser Asn Thr Tyr Leu His
 1 5 10 15
 15 <210> 186
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 20 <220>
 <223> mutant antibody L chain
 <400> 186
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Trp Asn Thr Tyr Leu His
 25 1 5 10 15
 <210> 187
 <211> 16
 <212> PRT
 30 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 187
 35 Arg Ser Ser Gln Ser Leu Val His Ser Asn Tyr Asn Thr Tyr Leu His
 1 5 10 15
 <210> 188
 <211> 16
 40 <212> PRT
 <213> Artificial Sequence
 <220>
 45 <223> mutant antibody L chain
 <400> 188
 Arg Ser Ser Gln Ser Leu Val His Ser Asn Arg Asn Thr Tyr Leu His
 1 5 10 15
 50 <210> 189
 <211> 16
 <212> PRT
 <213> Artificial Sequence
 55 <220>

<223> mutant antibody L chain

<400> 189

Arg Ser Ser Gln Ser Leu Val His Ser Asn Val Asn Thr Tyr Leu His

1 5 10 15

<210> 190

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 190

Arg Ser Ser Gln Ser Leu Val His Ser Asn Pro Asn Thr Tyr Leu His

1 5 10 15

<210> 191

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 191

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser

20 25 30

Asn Ala Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser

35 40 45

Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro

50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile

65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn

85 90 95

Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys

100 105 110

<210> 192

<211> 112

<212> PRT

<213> Artificial Sequence

<220>

<223> mutant antibody L chain

<400> 192

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Asp Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 10 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 15 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 193
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 193
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 30 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Glu Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 35 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 40 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 45 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 194
 <211> 112
 <212> PRT
 50 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 194
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 5 20 25 30
 Asn Phe Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 10 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 15 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 195
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 195
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 30 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn His Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 40 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 45 <210> 196
 <211> 112
 <212> PRT
 <213> Artificial Sequence
 50 <220>
 <223> mutant antibody L chain
 <400> 196
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Asn Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 10 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 15 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 197
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 197
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 30 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Thr Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 40 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 45 <210> 198
 <211> 112
 <212> PRT
 50 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 198
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Gln Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 10 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 15 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 199
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 199
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 30 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Ile Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 35 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 40 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 45 <210> 200
 <211> 112
 <212> PRT
 <213> Artificial Sequence
 50 <220>
 <223> mutant antibody L chain
 <400> 200
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Lys Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 10 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 15 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 201
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 201
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 30 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Leu Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 35 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 40 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 45 <210> 202
 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 50 <223> mutant antibody L chain
 <400> 202
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 5 20 25 30
 Asn Ser Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 10 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 15 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 203
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 203
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 30 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Trp Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 40 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 45 100 105 110
 <210> 204
 <211> 112
 <212> PRT
 50 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 204
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

EP 1 674 111 A1

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 5 Asn Tyr Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 10 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 15 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 <210> 205
 20 <211> 112
 <212> PRT
 <213> Artificial Sequence
 <220>
 25 <223> mutant antibody L chain
 <400> 205
 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 30 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Arg Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 35 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 40 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 45 <210> 206
 <211> 112
 <212> PRT
 50 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 206
 55 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly

1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 5 20 25 30
 Asn Val Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 10 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 15 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 20 <210> 207
 <211> 112
 <212> PRT
 25 <213> Artificial Sequence
 <220>
 <223> mutant antibody L chain
 <400> 207
 30 Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 35 Asn Pro Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 40 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 45 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Asn
 85 90 95
 Thr His Val Pro Pro Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 50

Claims

- 55 1. An antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 of any one of (1) - (12):

(1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 123, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 124, and CDR3 comprising the amino acid sequence set forth in SEQ ID

NO: 125;

(2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 109, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 110, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 111;

(3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 106, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 107, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 108;

(4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 132, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 133, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 134;

(5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 106, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 135, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 136;

(6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 126, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 127, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 128;

(7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 129, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 130, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 131;

(8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 103, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 104, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 105;

(9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 118, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 121, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 122;

(10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 115, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 116, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 117;

(11) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 112, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 113, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 114; or

(12) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 118, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 119, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 120.

2. An antibody comprising a light chain variable region having CDRs 1, 2 and 3 of any one of (1) - (13):

(1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 143, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;

(2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 143, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 145;

(3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 140, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 141, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 142;

(4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 167, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 168, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 169;

(5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 170, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 171;

(6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 159, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 160, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 161;

(7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 162, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 147, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 163;

(8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 164, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 165, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 166;

(9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 137, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 138, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 139;

(10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 155, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 156, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 157;

(11) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 149, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 150, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 151;

(12) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 146, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 147, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 148; or

(13) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 152, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 153, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 154.

3. An antibody selected from the group consisting of any one of (1) - (13):

(1) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 143, 144 and 158, respectively;

(2) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 109, 110 and 111, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 143, 144 and 145, respectively;

(3) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 106, 107 and 108, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 140, 141 and 142, respectively;

(4) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 132, 133 and 134, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 167, 168 and 169, respectively;

(5) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 106, 135 and 136, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 170, 144 and 171, respectively;

(6) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 126, 127 and 128, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 159, 160 and 161, respectively;

(7) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 129, 130 and 131, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 162, 147 and 163, respectively;

(8) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 129, 130 and 131, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 164, 165 and 166, respectively;

(9) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 103, 104 and 105, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 137, 138 and 139, respectively;

(10) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 118, 121 and 122, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 155, 156 and 157, respectively;

(11) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 115, 116 and 117, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 149, 150 and 151, respectively;

(12) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 112, 113 and 114, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 146, 147 and 148, respectively; and

(13) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid

sequence set forth in SEQ ID NO: 118, 119 and 120, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 152, 153 and 154, respectively.

4. An antibody having a heavy chain variable region of any one of (1) - (7):

- (1) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84;
- (2) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85;
- (3) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86;
- (4) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87;
- (5) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 88;
- (6) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 89; or
- (7) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 90.

5. An antibody having a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92.

6. An antibody selected from the group consisting of the antibody of any one of (1) - (7):

- (1) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (2) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (3) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (4) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (5) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 88 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92;
- (6) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 89 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92; and
- (7) an antibody comprising a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 90 and a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 92.

7. An antibody having an activity equivalent to the activity of the antibody of any one of Claims 1-6, wherein one or more amino acid residues are substituted, deleted or added and/or inserted from the amino acid sequences set forth in any one of Claims 1-6.

8. The antibody as claimed in any one of Claims 1-7 which is a humanized antibody.

9. A humanized antibody capable of binding to glypican 3.

10. An antibody capable of binding to a peptide consisting of the sequence of the amino acid residues 524 - 563 of glypican 3.

11. An antibody capable of binding to a peptide consisting of the sequence of the amino acid residues 537 - 563 of glypican 3.

12. The antibody as claimed in Claim 10 or 11 which does not bind to a peptide consisting of the sequence of the amino acid residues 550 - 563 of glypican 3.

13. An antibody capable of binding to a peptide consisting of the sequence of the amino acid residues 544 - 553 of glypican 3.

14. An antibody capable of binding to a peptide consisting of the sequence of the amino acid residues 546 - 551 of glypican 3.

15. The antibody as claimed in any one of Claims 9-14 which is a humanized antibody.

16. An antibody capable of binding to an epitope to which a second antibody is capable of binding, wherein said second

antibody comprises a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 143, 144 and 158, respectively.

- 5 17. An antibody capable of binding to glypican 3 which has a high CDC activity against a cell expressing glypican 3.
18. An antibody capable of binding to glypican 3 which has a high ADCC activity against a cell expressing glypican 3.
19. A polynucleotide coding for a heavy chain variable region or a light chain variable region of the antibody as claimed
10 in any one of Claims 1-18.
20. The polynucleotide as claimed in Claim 18 having the sequence set forth in SEQ ID NOs: 11-21, 33-43, 55-59, 65-70 and 77-83.
- 15 21. A cell-growth inhibitor comprising as an active ingredient the antibody as claimed in any one of Claims 1-18.
22. An anticancer agent comprising as an active ingredient the antibody as claimed in any one of Claims 1-18.
23. The anticancer agent as claimed in Claim 22 for treatment of hepatoma.
- 20 24. A peptide comprising the amino acid sequence of the amino acid residues 524 - 563 of glypican 3.
25. A peptide comprising the amino acid sequence of the amino acid residues 537 - 563 of glypican 3.
- 25 26. A peptide comprising the amino acid sequence of the amino acid residues 544 - 553 of glypican 3.
27. A peptide comprising the amino acid sequence of the amino acid residues 546 - 551 of glypican 3.
28. An antibody comprising a light chain variable region having CDRs 1, 2 and 3 of any one of (1) - (15):
30 (1) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 174, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
35 (2) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 175, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
(3) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 176, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
40 (4) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 177, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
(5) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 178, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
45 (6) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 179, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
(7) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 180, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
50 (8) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 181, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
55 (9) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 182, CDR2 comprising the amino acid sequence set forth in SEQ ID NO: 144, and CDR3 comprising the amino acid sequence set forth in SEQ ID NO: 158;
(10) CDR1 comprising the amino acid sequence set forth in SEQ ID NO: 183, CDR2 comprising the amino acid

CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 186, 144 and 158, respectively;
 (14) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid
 sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having
 CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 187, 144 and 158, respectively; and
 (15) an antibody comprising a heavy chain variable region having CDRs 1, 2 and 3 comprising the amino acid
 sequence set forth in SEQ ID NO: 123, 124 and 125, respectively, and a light chain variable region having
 CDRs 1, 2 and 3 comprising the amino acid sequence set forth in SEQ ID NO: 188, 144 and 158, respectively.

30. An antibody having a light chain variable region selected from (1) - (15):

- (1) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 191;
- (2) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 192;
- (3) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193;
- (4) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 194;
- (5) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 195;
- (6) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 196;
- (7) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 197;
- (8) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 198;
- (9) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 199;
- (10) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 200;
- (11) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 201;
- (12) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 202;
- (13) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 203;
- (14) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 204; and
- (15) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 205.

31. An antibody having a light chain variable region selected from the group consisting of (1) - (15):

- (1) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 191;
- (2) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 192;
- (3) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 193;
- (4) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 194;
- (5) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 195;
- (6) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 196;
- (7) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 197;
- (8) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 198;
- (9) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 199;
- (10) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 200;
- (11) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 201;
- (12) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 202;
- (13) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 203;
- (14) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 204; and
- (15) a light chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 205; and a heavy
 chain variable region selected from the group consisting of (1) - (7):

- (1) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 84;
- (2) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 85;
- (3) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 86;
- (4) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 87;
- (5) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 88;
- (6) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 89; and
- (7) a heavy chain variable region comprising the amino acid sequence set forth in SEQ ID NO: 90.

32. The antibody as claimed in any one of Claims 28 - 31 which is a human antibody.




FIG.1

Mab	Inhibition of the binding of indicated biotin-Mab to sGPC3 core (%)								Topographical epitope
	M3C11	M1E7	M11F1	M6B1	M18D4	M5B9	M10D2	L9G11	
M3C11	<u>96.8</u>	<u>96.2</u>	12.0	11.2	10.3	9.1	39.9	33.1	a
M13B3	<u>71.3</u>	<u>95.7</u>	15.6	9.3	2.7	-1.8	22.3	24.9	
M1E7	26.0	<u>83.9</u>	14.3	7.7	9.0	6.4	14.8	38.0	
M3B8	1.8	28.4	<u>97.4</u>	15.2	12.6	3.6	29.0	36.0	b
M11F1	2.0	3.9	<u>81.8</u>	5.6	-1.5	5.4	24.6	20.9	
M19B11	-4.1	-9.4	3.1	<u>87.6</u>	44.1	31.0	14.2	-9.8	c
M6B1	-5.7	-2.7	13.9	<u>85.6</u>	<u>51.9</u>	44.4	6.7	1.1	
M18D4	0.2	0.5	-0.3	38.6	<u>89.0</u>	<u>89.4</u>	19.1	6.7	
M5B9	0.1	2.7	5.3	23.2	<u>77.3</u>	<u>77.3</u>	13.4	10.3	
M10D2	-7.7	0.6	7.8	-5.9	-21.6	-10.3	<u>79.2</u>	7.2	d
L9G11	2.1	-4.8	6.0	10.1	15.7	6.4	1.2	<u>92.2</u>	e

FIG.2




FIG.3




FIG.4




FIG.5




FIG.6




FIG.7




FIG.8




FIG.9




FIG.10




FIG.11




FIG.12

	GC33 WB
GC-4	GNSQQATPKDNEISTFHNLGNVHSP LK
GC-5	STFHNLGNVHSP LK
GC-6	GNSQQATPKDNEIS
GC-7	GNSQQATP
GC-8	QQA TP KDN
GC-9	TP K DNE IS
GC-11	ATPKDNEIST
GC-12	PKDNEISTFH
GC-13	DNEISTFHN L
GC-14	EISTFHN L GN

FIG. 13




FIG.14




FIG.15

Clone ID	Isotype	ELISA	BIACORE			EPTOPE		FACS				Immuno-precipitation		
		EC50 (nM)	ka (1/Ms /10 ⁵)	kd (1/s x10 ⁵)	KD (nM)	Competitive ELISA	western blotting	CHO/ human GPC3	HepG2	HuH-7	CHO			
M3C11	IgG1	0.12	2.5	6.5	0.3	a	GC-2	83	59	9	-	Δ		
M13B3	IgG1	0.25	N.D.	N.D.	N.D.			N.D.	N.D.	N.D.	-	N.D.	-	N.D.
M1E7	IgG1	0.96	1.5	42.7	2.8			28	16	5	-	-	x	
M3B8	IgG1	0.56	2.7	467.0	17.6	b	GC-4	12	6	3	-	x		
M11F1	IgG2b	2.17	1.4	3.5	0.2			9	4	2	-	-	x	
L9G11	IgG1	0.62	1.3	140.0	10.7	c	N	29	26	4	-	○		
M19B11	IgG1	0.18	1.5	17.1	1.1			39	44	3	-	-	○	
M6B1	IgG1	5.51	N.D.	N.D.	N.D.	37		34	3	-	-	○		
M18D4	IgG1	0.85	0.8	49.6	6.2	22		20	2	-	-	○		
M5B9	IgG1	1.13	0.6	139.0	23.5	25		23	2	-	-	○		
M10D2	IgG1	1.04	1.1	49.6	4.7	d		20	29	3	-	-	○	

FIG.16

Clone ID	Isotype	ELISA	EPITOPE		FACS			
		EC50 (nM)	Competitive ELISA	Western blotting	CHO/human GPC3	HepG2	HuH-7	CHO
GC199	IgG1	0.10	b	GC-4	15.5	12.2	2.4	-
GC202	IgG1	0.10			6.0	0.9	1.4	-
GC33	IgG2a	0.24			82.7	52.0	8.4	-
GC179	IgG2b	5.61	f	GC-3	7.1	6.5	3.8	-
GC194	IgG1	3.83			5.0	7.9	1.5	-

FIG.17




FIG.18




FIG.19




FIG.20

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP2005/013103

A. CLASSIFICATION OF SUBJECT MATTER

C07K16/18(2006.01), C12N15/09(2006.01), A61K39/395(2006.01),
A61P1/16(2006.01), A61P35/00(2006.01), C07K14/47(2006.01),
C12P21/08(2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K16/18(2006.01), C12N15/09(2006.01), A61K39/395(2006.01),
A61P1/16(2006.01), A61P35/00(2006.01), C07K14/47(2006.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS/WPI(DIALOG), CA(STN), REGISTRY(STN), GenBank/EMBL/DDBJ/GeneSeq,
SwissProt/PIR/GeneSeq

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2004/022739 A1 (Chugai Pharmaceutical Co., Ltd.), 18 March, 2004 (18.03.04), & AU 2003261941 A1 & EP 1541680 A1	1-32
X	WO 2004/023145 A1 (Kabushiki Kaisha Peruseusu Proteomikusu), 18 March, 2004 (18.03.04), & AU 2002330482 A1	1-32
X	WO 204/038420 A1 (Kabushiki Kaisha Peruseusu Proteomikusu), 06 May, 2004 (06.05.04), & AU 2003261943 A1 & EP 1548442 A1	1-32

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
11 October, 2005 (11.10.05)

Date of mailing of the international search report
25 October, 2005 (25.10.05)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.